



Credit: 2 PDH

Course Title:

Is Insulin Resistance Work Related

Approved for Credit in All 50 States

Visit epdhonline.com for state specific information including Ohio's required timing feature.

3 Easy Steps to Complete the Course:

1. Read the Course PDF
2. Purchase the Course Online & Take the Final Exam
3. Print Your Certificate

Is Insulin Resistance Work Related?

Marina Oțelea

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/66333>

Abstract

Incidence of insulin resistance continues to grow, becoming a major public health concern worldwide. Besides the classical risk factors (obesity, unhealthy nutrition and lack of exercise), extensive research about some occupational hazards supports their association with insulin resistance and metabolic syndrome. On the one hand, the classical risk factors for insulin resistance are augmented by the changes of the working conditions: the occupational level of physical activity has a tendency to decrease, reducing its contribution to the overall level of physical activity and favouring a sedentary lifestyle and the occupational stress became the second most common work-related health issue, contributing to the increase of the maladaptive habits, such as unhealthy nutrition. On the other hand, some insulin resistance risk factors are primarily occupational hazards: desynchronization of the circadian rhythm and sleep disruption during the night shifts, workplace air pollution (particles, solvents), heavy metals (arsenic, mercury) or persistent organic pollutants exposure. Meantime, workplaces are excellent settings for health-promotion programmes and metabolic risk reduction, if there is managerial commitment and support. Therefore, assessment of the risk, screening and workplace intervention programmes to reduce insulin resistance incidence should be included in the occupational health service provision.

Keywords: insulin resistance, occupational stress, sleep desynchronization, night shifts, sleep duration, indoor air pollution, persistent organic pollutants arsenic, mercury

1. Introduction

Insulin resistance (IR) is a complex pathophysiological state characterized by reduced target cells responsiveness to insulin. The variety of its manifestations correspond to the complexity of insulin actions in specific tissues; the IR state reduces glucose uptake in muscle and other insulin-dependent tissues, increases lipolysis and fatty acids delivery from adipose tissue, increases hepatic glucose output and sodium retention, impairs uric acid metabolism,

enhances vascular endothelium vasoconstriction and inflammation—promoting atherosclerosis and induces a systemic low grade metabolic inflammation. Two are the major clinical consequences of the IR: the cardio-vascular disease and diabetes. There are also significant links between IR and other clinical conditions such as non-alcoholic fatty liver disease, polycystic ovary syndrome and colorectal cancer. Cardiovascular diseases are already the leading cause of death in industrialized countries [1] and diabetes prevalence has doubled between 1980 and 2014. Diabetes is estimated to become the seventh leading cause of death in 2030 [2]. Taken together, all medical conditions related to IR are responsible for a high percentage of the total mortality rate. Therefore, prevention and early diagnosis is a major public health topic.

For many years, IR has been considered a complication of obesity but a non-obese metabolic obese profile has been identified in the past few years, as up to 15–30% of the IR subjects are in the normal range value of the BMI. The complex pathogeny of IR has been explained by complementary or synergic interactions between genetic and epigenetic factors, gut microbiota and environmental factors. As the environmental factors are the modifiable ones, the intervention, including workplace interventions, became nowadays a major health prevention direction.

Insulin resistance is difficult to measure in clinical practice; therefore, surrogates are used for its assessment such as basal insulinemia, the homeostasis assessment index (HOMA-IR), glucose ratio to insulin and the glucose tolerance test. Clinical markers for IR are lipid metabolism impairment (increased triglycerides, reduced HDL cholesterol and increased LDL cholesterol), plasma uric acid increase, renal disease markers (albuminuria) and cardiovascular effects (hypertension and cardiac remodelling). As already mentioned, obesity, particularly the visceral deposition of the adipose tissue, is a risk factor. In different configurations, according to different scientific organizations, these factors define the metabolic syndrome, a more suitable clinical diagnostic. Although I fundamentally agree with those considering that metabolic syndrome is an ‘artificial syndrome’—in fact is just a cluster of risk factors—and that diagnostic procedures should be developed for an easier clinical diagnostic of the IR syndrome, this article will have to rely on data accumulated under the metabolic syndrome entity, as it is much more frequently used in the medical literature.

Nutrition, lack of physical exercise are not occupational hazards that are traditionally IR risk factors. In recent research, there is more and more evidence accumulated on how particular occupational risks are involved in IR and this makes the subject of increasing interest for occupational physicians.

2. Epidemiological arguments

An impressive number of studies have emphasized that risk factors for cardiovascular and diabetes are more prevalent in low-educated, low-income population [3–8], particularly related to the high incidence of the negative health habits (smoking, drinking, high fat or low fruit diet [8]). Being a member of a low social class during childhood [9] also favoured unhealthy-related behaviours: low level of sport practice, high fat diet, emotional stress and material unfavourable conditions. In different cohorts, significant differences between men

and women [6, 7] were found, which could not be fully explained by sexual level of hormones. Marital status [5] in men appeared to have different influences in developing countries compared to the developed ones: a high risk factor in developing countries and a low risk factor in western countries [4]. The high prevalence in developing countries has been explained by cultural habits such as accessibility to eat out or practice outdoor physical activities and less concern about gaining weight. In women, marital status in western countries is associated with an increased risk.

In the early years of socio-economic development, the changing habits represent a risk factor for IR [10, 11]. The theory of epidemiological transition states that, under development conditions, there is a switch between mortality dominated by infectious diseases to mortality related to degenerative and man-made diseases [12]; conditions related to IR fill in this category, reflecting changes in nutrition and transition from physical activity to sedentariness [11]. In these countries, at the beginning of the development era, the high income class has higher risk [5], as it has access to more resources, moves to more sedentary lifestyle and tries to compensate all childhood economical disadvantages and frustrations accumulated.

Even if there are differences according to the socio-economic development period the country is passing by, there are also many similarities. For example, a comparison between Taiwan and US revealed interesting similar inflammation scores (serum IL-6, CRP, fibrinogen, sICAM-1, sE-selectin and sIL-6R), cardiovascular risk factors (systolic and diastolic blood pressure, total cholesterol, HDL cholesterol, triglycerides, glycosylated haemoglobin, BMI, waist-hip ratio and resting pulse) and hypothalamic-pituitary-adrenal axis and sympathetic nervous system function (epinephrine, norepinephrine, DHEAS and urinary cortisol) in relation to the socio-economic status [13].

Occupation and employment status are major determinants of the economical welfare and unemployment has been linked to metabolic risk in both developed and in under development countries [4, 5, 14]. There is difficult to find common grounds for data reported from different countries, as occupations are differently coded and registered: an US countrywide survey found maximum incidence of risk factors in the farm industry (operators, managers and supervisors), in food preparation and food services, motor vehicle operators and among administrative staff (secretaries, stenographers and typists), and the lowest in professional specialties (particularly in writers, artists, entertainers and athletes) [15]. Several studies have classified occupations in manual (blue collars) and non-manual (white collars, office workers) activities and found either increased or a decreased incidence of metabolic syndrome components in manual workers [16, 17]. These differences were most probably related to the confounders not included in these studies, such as nutrition, smoking, alcohol and level of occupational physical activity.

There are many limitations of these epidemiological studies for a direct application in the occupational physician practice, although the prevalence and social risk factors are important. They give a broader perspective for any intervention proposed. The epidemiological transitions reflect the major trends and do not provide specific information by industry. As far as working conditions are concerned in the transitional period, the main negative factors are the unbalanced working hours to leisure activities, the increased number of jobs with sedentary activity and the increased amount of perceived stress.

3. Occupational hazards

3.1. Sedentary lifestyle

The link between inactivity and insulin resistance has been documented for many years and extensive experimental and clinical research have clarified this relationship in detail. Currently, an inactive person is defined by having less than 30 min/day of at least moderate level of activity, either as occupational or as leisure time activity. As the occupational activities become less physical demanding, it would be reasonable to compensate by more physical activities such as sports practice, gardening and brisk walking [18] instead of increasing the number of hours of TV watching or computer interactions, as we see as a trend nowadays [19]. It is estimated that only 20% of currently industrial jobs demand an average level of physical activity, with a reduction of more than 100 kcal/day in energy expenditure since 1960 [18].

Benefits of exercise in metabolic regulation: Glucose is transported into muscle cell by facilitated diffusion through GLUT4 and GLUT1 channels; GLUT4 is the major transporter. In resting conditions, GLUT4 is located intracellularly and its traffic to the membrane is insulin mediated. Inside the muscular cell, glucose is phosphorylated to gluco-6-phosphate (G-6P), a compound that cannot transfer outside the cell. During a rest, it is stored as glycogen and during exercise it takes the glycolytic pathway. During exercise, the blood flow increases, thus increasing the amount of glucose available; this creates a gradient for glucose to enter the cell. As cells utilize energy for sustaining muscular contraction, the ratio of AMP/ATP increases, activating the AMP-activated protein kinase (AMPK). This kinase is able to enhance the transport of GLUT4 to the membrane in an independent insulin manner. AMPK activity is also increased by leptin, through indirect hypothalamic action and adiponectin, by a direct action. In obesity, resistance to leptin diminishes AMPK activity [20]. Exercise increases GLUT4 transport and fatty acid oxidation, improving insulin sensitivity. Acute changes depend on the energy balance: if this is positive, the carbohydrate oxidation will prevail and if it is negative, the fatty acid oxidation will be enhanced [20]. In order to reduce fatty mass, a negative balance is also requested. The type of exercise also influences the energy substrate utilization: endurance training increases fatty acid oxidation while resistance training, increases glucose uptake and utilization; most probably, this is due to the increase in muscular mass [21]. Lipoprotein lipase increases after training and maintains a higher level afterwards; the higher the level of exercise, the longer will be the duration of the high level of lipoprotein lipase and the preventive action. Chronic training contributes to increase the number of myofibrils and their oxidative capacity. Training has also a role in restoring mitochondrial content in ageing populations or in different pathological conditions [22–25] and in activating satellite cells [26]. The mitogenesis effect increased with 25% when calorie restriction was associated [27]. In previous sedentary obese men and women, training increased the mitochondrial electron transport and decreased IR.

Exercise partially counteracts other risk factors, such as a fatty diet. High lipid intake increases the fatty acids content in muscular cells, β -oxidation, acetylCoA level, NADH/NAD ratio (slowing of tricarboxylic acid cycle) and leads to citrate accumulation. Citrate inhibits muscle hexokinase and reduces the formation of gluco-6-phosphate; the glucose traffic through

GLUT4 channels is, therefore, reduced, contributing to chronic hyperglycaemia. This process favours fatty acid utilization and counteracts the insulin metabolic switch. A vicious pathological cycle is created between decreased sensitivity of GLUT 4 and reduced uptake of glucose. Some benefits of exercise are explained by disrupting this cycle. Training even reversed the other negative risk factors, such as obesity influence or small size at birth [28].

Relationship between sedentary lifestyle and obesity: Sedentary lifestyle infers a low energy expenditure that should be adjusted by a balanced reduction in food intake, but food intake is driven by many other factors compared to expenditure in abundant, accessible nutrient societies. In such context, calorie intake becomes easy and higher than requested for thermo-regulation and metabolic purposes. In a strict mathematical formula, this will lead to increase in body weight. High calorie intake is a consequence of high fat and/or high-carbohydrate diets. Randle's hypothesis of fatty acid toxicity [29] is a classical explanation of IR induction by excessive fatty diets. High circulating level of fatty acids inactivates insulin cellular signal and impairs the switching of the metabolism from fatty acids to glucose, with progressive accumulation of intra-cytoplasmic lipids in insulin sensitive cells (including β -cells); the lipid accumulation, mainly of ceramides, induces apoptosis. In muscular cells, the increased level of fatty acids inhibits phosphorylation of insulin receptor substrate 1 and activation of phosphoinositol 3 kinase, with a decrease in glycogen storage. High carbohydrate diets, particularly of those with high glycaemic index, desensitise glucose receptors and increase oxidative stress on β -cells that ultimately lead to insulin depletion. High fructose intake feeds the gluconeogenesis and the lipogenic pathways [30] in hepatic cells and by depleting the cellular ATP content and gives to the brain a signal of energy deprivation, favouring food seeking behaviour. Increased hepatic glucose output maintains a hyper-insulinemic state with higher fat storage in adipose tissue.

Sedentary life decreases the proportion between the fat mass and the free fat mass. According to the energy balance concept, it would be expected, as the total body mass increases, that the total energy consumption increases, from the basal metabolism increase. The energy expenditure is dominated (with exception for highly competitive professional athletes) by the resting metabolic rate and the variation in size of the free fat mass that explains 60–80% of the variation in basal metabolism [31], with an approximated 5:1 ratio between free fat mass and fat mass [32]. Energy balance is a remarkable concept, but cannot fully explain the complexity of the metabolism regulation and of the metabolic flexibility, governed by insulin. In sedentary adult with positive energy balance, the proportion of fatty mass increases gradually; even if the adjusted increase of the basal metabolic rate would be proportionate with the total mass, this increase will not compensate the level of calories intake, because fat mass has lower basal metabolism than skeletal muscle and fat will continue to accumulate. Obesity, as defined by a quantitative increase of fat tissue or as proportion between fat mass and fat free mass [33], is a direct consequence. The abnormal proportion and not the BMI is, in fact, related to the metabolic risk factors. This explains, at least partially, the cardio-metabolic risk of the non-obese metabolically obese population. This increase in quantity is also associated with qualitative impairments, a phenomenon described as adiposopathy or 'sick fat' [34]. The increased level of leptin and of other inflammatory cytokines, the decreased level of adiponectin, switch of the macrophages to the M1 pattern and to a pro-inflammatory status and lower capacity of

oxidizing fat are the main characteristics of the metabolic pattern of adiposopathy. Muscle is the main consumer of glucose and fatty acids and by that, the major contributor to the lowering of glucose and of the fatty acid plasma levels; if free fat mass (mainly muscle mass) is reduced due to the sedentary lifestyle, the risk of adiposopathy increases.

Occupational level of physical activity has a general tendency to decrease. This reduction in contribution to the overall level of physical activity explains that a significant IR risk was associated with leisure time sedentary behaviour but not with occupational activity in general population [35]. However, when looking more closely, it becomes obvious that several professions are at higher risk: for example, VDU operators in call centres [36] or bus drivers [37] had a higher OR for metabolic syndrome, after confounders such as smoking or leisure time activity were excluded. Comparison of sedentary lifestyle has been highly suggestive between different professions and even among same profession that has specificities according to national regulations: for example, police officers and administrative clerks have the higher risk than fire fighters in states where training programmes are strictly supervised but not in voluntary forces (US) [38].

During work, the sedentary activity can be more sustained (more than 30 min) than during non-working hours and this has also an impact [39]. Beneficial effects in adiposity measures, triglycerides and 2-h plasma glucose were obtained even in sedentary persons with more interruptions in sedentary time [40].

Sitting for more than 1 h reduces the lipoprotein lipase activity [41] and the non-exercise activity thermogenesis [42]. Adjustable workstations might be a solution for prolonged sitting activity. In a recent systematic review, the majority of prospective studies found that occupational sitting was associated with a higher risk of diabetes and mortality and to some degree of cardiovascular risk. These apparent conflicting results between the two consequences of the IR could be related to the heterogeneity of study designs and to the inconsistency and sufficiency of data recorded on overall physical activity. Possible other confounders are diet, tobacco, alcohol and high energy food consumption, as sitting for long hours increases the chance of negative habits to develop. When the total physical activity was considered, the association between sitting and metabolic risk lost the statistical significance [43].

Clinical approach: Clinical evaluation of the physical activity is rather easy and should be part of the routine examination. BMI is also very frequently recorded and, even if it has the above-mentioned limitations, it has an orientation value. In order to be more specific about the metabolic risk, abdominal circumference could be added to the physical examination, particularly to the non-obese persons, defined by the BMI index. More accurate data about fat mass proportion are obtained by bioelectrical impedance analysis, but they are not used in routine clinical examinations.

As high physical activity has numerous benefits, it should be encouraged to everyone, according to its age, health status and preferences. There is enough evidence that even small increases, if they are sustained and become part of the routine, have high medium-term impact as primary or secondary preventive measures. If sitting is the main problem, an ergonomic intervention could be beneficial.

3.2. Sleep and night shifts

Sleep duration and sleep architecture are important indicators of individual health. When normal sleep is impaired, alertness, memory, learning, thermoregulation, respiratory, metabolic and cardio-vascular regulation are also disturbed. This broad spectrum of effects is one of the consequences of the complex connections between the hypothalamic clock and different nuclei in the central nervous system [44] and of the interaction between the central and the peripheral circadian regulators. Assessing sleep duration, sleep-wake cycle and sleep disorders in the occupational medical practice is very important, because they influence significantly the fitness for work particularly in high risk jobs for accidents (professional driving, traffic controllers, working at height, etc.).

Normal sleep regulation: Sleep onset and duration are the core elements of sleep-wake regulation. Sleep-wake regulation is currently explained by two biological models: the sleep homeostasis and the circadian clock. Sleep homeostasis refers primarily to the variation of sleep pressure, while the circadian clock concerns the activity of the hypothalamic pacemaker, acting independently of the prior sleep period [45]. Although interrelated, their regulation pathways are different: sleep pressure is mainly dependent on the existing sleep debt and is the physiological contributor to the recuperation of this debt. If during the previous day the sleep duration was shorter, the sleep homeostasis will intervene to correct the deficit.

The pacemaker of the clock automatism is genetically set to a circadian rhythm and is controlled by the environmental factors. The circadian period (the interval that covers a full cycle of sleep-wake, not influenced by external constraints) is the *tau*; *tau* varies in healthy individuals between 24 and 25 h and reflects the automatism of the suprachiasmatic nuclei processes (is circadian clock dependent). In healthy adult individuals, periodicity of sleep duration waxes and wanes ranges broadly (around 2–18 days), indicating that for a specific individual the number of hours slept varies in consecutive days, having a repetitive pattern of duration. For example, for the 5 days pattern, the number of hours slept in day 1 is the same as the one in day 6; between day 1 and day 6 there is an increase and then a decrease in duration of sleep. This inter-individual variation is determined by the different build up sleep pressure in healthy individuals and is dependent primarily to the sleep homeostasis. This variance will impact on different ways the adaptation to night shifts and to longer working hours of different individuals.

Under a certain extent, the clock is self-regulated by its genomic functionality. The activation of the hypothalamic neurons starts with the transcription of the *Clock* and *Bmal 1* genes that have circadian variations, related to day-night transitions; their products form a heterodimer (CLOCK-BMAL1) that binds to the promotion site of *Period 2* (*Per2*), *Cryptochrome* (*Cry*). The cytoplasmic levels of PER2 and CRY regulate the transcription induced by CLOCK-BMAL1 heterodimer through a negative feedback loop. Besides this auto-regulated feedback loop, *Clock* and *Bmal1* are also influenced by a secondary control loop, acting on two nuclear receptors, the *Retinoid-related orphan receptor* (*Ror*) and the *Rev-Erb receptor*. While *Ror* is a transcription activator, *Rev-Erb* is an inhibitor one [46]. PER and CRY reach a maximum at the normal transition from day to night maintaining the negative feedback loop of the clock and contributing to the sleep onset. BMAL1 reaches the maximum at the transition from day to night, at

waking time. In order to generate the level of BMAL1 requested for waking, ROR and REV ERB maxima occur between that of BMAL1 and PER-CRY.

Dark is the physiological stimulus of melatonin, although melatonin increases during the evening even in blind persons [47]. While its major role is to induce sleep, melatonin has also other functions such as heat loss or glutathione synthesis inducer [48]. It also influences the immune response and the glucose metabolism.

Sleep cycles have normal variation during the night: in the first part of the night, there are more non-rapid eye movement (NREM) periods than in the last part; stage 1 of sleep diminishes (in percentage time of the total cycle) in the second part of the night and stage 1 is no more present after the first 4 h of night sleep. During nightshifts or sleep interruptions, the normal sleep architecture is modified and explains the tiredness people feel, after a working night, even after an equal number of hours of recovery sleep.

Normal sleep duration: A large European study revealed significant differences between countries, related mainly to cultural habits than to environmental ones (sunlight or temperature variation) as winter and summer sleep duration showed no significant variation by country [49]. On average, adults sleep 7–7.5 h/night [50]. In adults, adjusted prevalence of short sleepers (<5–6 h/night) or long sleepers (>10 h/night) is 0.52 and 0.64%, respectively [51]. Duration of sleep is influenced by genetic, personality traits and social habits. Long sleepers had higher neuroticism scores and lower education levels than short sleepers [51, 52]. Short sleepers are more frequent in adults that have not sleep alone in their childhood, as toddlers or pre-schoolers [52]. Familial nocturnal short sleepers have been identified in a mutation on the gene *DEC2*: in persons carrying this mutation, the whole duration of sleep in 24 h is less than 6 h [53]. Shorter sleeping time was associated with the total time spent sitting and with increased screen time in a large cross-sectional European survey [54], linking this risk factor with sedentary lifestyle and with a long time of computer interaction.

Metabolic effects related to sleep regulation: Hypothalamic clock does not regulate just the sleep-wake status; it is the central driver for the peripheral clocks correspondents in colon, stomach, pancreas, liver, adipose tissue and muscles: through this integration, certain metabolic pathways have circadian fluctuation. It is estimated that 10–15% of the gene transcripts have rhythmically 24-h cycles; besides the genes expression, translational changes have also circadian fluctuations. There is a normal variation of the peak transcripts during day (e.g. proteins involved in inflammation and hormonal response) or night (e.g. proteins involved in ribosomal and epigenetic processes such as methyltransferases) [55]. In twin-based genetic studies, heritability is found responsible for 33–55% of the sleep duration; interestingly, heritability influence increases when twins are living together for a longer period in their life [56].

Besides the central clock influence, there is also an intrinsic circadian variance of the metabolic genes, as response of the *Clock/Bmal1* or *Per 2* variation of expression. All hormones involved in glucose regulation (cortisol, catecholamine, glucagon and insulin) have circadian peaks. There is significant increase in insulin sensitivity during the early nocturnal sleep and a returning to the pre-sleep values during the late phase of the sleep [57]. In animal studies, melatonin decreases insulin and increases glucagon secretion in pancreatic cells; high insulin

levels, as reached in type II diabetes, decrease the melatonin level [58]. In humans, several genetic polymorphisms of the melatonin gene are risk factors for IR. A later rise of the morning melatonin peak is encountered in persons with the GG allele of the melatonin gene and this time corresponds to the usual timing for food intake. The later than normal melatonin raise deregulates glucose metabolism, as it counteracts the nutrient-related signals transmitted to the pancreas cells. Other genetic variants, as the *MTNR1B* gene variant, increase the number of melatonin receptors expressed by β -cells with negative impact on the insulin secretion. Subjects carrying this allele, receiving 4 mg melatonin/day, had, after 3 months of treatment, significantly less insulin secretion and were more IR than controls [59]. Possible future testing of these genomic characteristics will allow occupational physicians to distinguish the persons that will have more difficulties in adapting to working in shifts and those that are at risk of developing metabolism disturbances, including IR.

Negative influences of work schedule on insulin resistance

1. *Short sleep duration.* Short-term sleep loss could be perceived as a metabolic stressor: it increases metabolism but also the anti-oxidant capacity [60]. These adaptations are surpassed in sleep deprivation, when major disturbances in hormonal regulation occur. Glucagon and cortisol are lower after a night of sleep deprivation [61]; there is also a lower response in adrenaline to hypoglycemia, explaining both the hunger and the fatigue syndrome in the day after. The day after sleep deprivation is characterized by both autonomic symptoms (anxiety, sweating, irritability, hunger) and glycopenic symptoms: dizziness, blurred vision and difficulty in thinking. After four nights of sleep deficit, the insulin signal pathway was reduced by 30% in adipocytes [62]. Cortisol levels increase just after one night of sleep deprivation. Even if the glucose tolerance tested by I.V. glucose tolerance test was similar with the reference group, the short sleepers had a reduction by 40% of the insulin sensitivity, both in the first and in the second phases of response [63]. Another strong argument of the influence of sleep on the IR is the reduction of HOMA-IR after an increase of amount of time slept by 45 min/night, a rather cost-effective intervention for prevention of IR [64].

The work time arrangements are very important for maintaining a physiological recovering sleep. Increased risk for metabolic syndrome was not constantly found in relation to working in shifts. However, alterations in glucose and lipid metabolism, observed in experimental settings [65, 66] of night shifts combined with a total reduction on sleeping hours (less than 5 h/day) and adverse nutritional habits [67] significantly increase the risk.

2. *Desynchronization of the circadian rhythm.* Another consequence of working in shifts is the shifted circadian rhythm (desynchronization, misalignment). The desynchronization increases the effects of sleep deprivation; the insulin sensitivity is twice reduced if restriction is combined with 8.5 sleep delay (circadian misalignment). After 10 days of longer day duration (duration of 28 h) a significant increase of glucose, decrease of leptin, reversed cortisol cycle and an increase of mean arterial pressure [68] have been recorded. A decrease in insulin sensitivity and leptin secretion are also common [69].

The metabolic effects of the desynchronization on the circadian rhythm have been explained by the conflict between the disturbed circadian variations of the peripheral clocks

and the feeding stimulus. In general, peripheral clock shifts at a slower rate than the central clock. The circadian variation of different peripheral factors involved in metabolism regulation such as adipokines [70], pancreatic secretions, PPAR γ PGC1 α is supported by numerous experiments. Besides this circadian variation, their expression is also triggered by the nutrition intake. For example, leptin plasma levels depend on meal time; irregular time of nutrient intake is a risk factor for the leptin resistance, a common finding in IR [71]. Pancreatic cell possesses self-sustained circadian genes and protein oscillations of the transcription factors clock bmal1 [72]. PPAR γ , a regulator of adipocyte differentiation and secretion and of insulin signal transmission, has circadian oscillations [73] but it is also activated by fatty acids. PGC1 α coordinates genes in lipid and glucose metabolism, the PAI-1 promoter and the thrombocytes activation [30]. It is stimulated, in heart and liver, by fasting [74]. What are the specific interferences between the intrinsic oscillation of expression of these proteins and the environmental factors (mainly nutrition and light) are currently a subject of intense research.

Social and psychological factors are other important modifiers of sleep homeostasis and even of the pacemaker control, with specific markers at the cellular level. For example, in forced-desynchrony experiments in which the sleep-wake cycle and the associated fasting-feeding and dark-dim light cycles are scheduled at a longer *tau* than 24 h, disrupted transcription of the clock genes expression, epigenetic modifications of chromatin, impairment of the RNA polymerase, modification of the regulation of transcription and translation of heat shock proteins and temperature-sensitive RNA binding proteins (CIRBP and RBM3) have been recorded [55].

The higher weight gain of the night eating syndrome is another consequence of the desynchronization of the circadian rhythm. A plausible biological explanation is that the nutrient-induced stimulus is acting out of the phase of the physiologic oscillations of these proteins. If, for example, melatonin secretion is delayed, there will be a decrease time between wake time and circadian melatonin offset phase that will increase the eating behaviour. The modification of appetite could also be related to changes in ghrelin and leptin levels in short sleepers, significantly increasing BMI [75]. Changes in preference for salty or high carbohydrate foods, irregular meals and snaking behaviours [76] have also been reported.

3. *Sleep disruption* is well studied particularly in sleep apnea, but in this specific sleeping disorder the intermittent hypoxia was the core mechanism for the explanation of the majority of the cardio-metabolic influences.

During the night shift, sleep disruption cannot be avoided. It certainly affects the quality of sleep and the sleep architecture (the proportion between REM and NREM cycles and their duration). Sleep (particularly REM cycles) activates transfer from the short- to long-term memory and allows, from the occupational perspective, to integrate new tasks of the job; therefore, time to recovery is an important factor influencing the stress level and the employees' productivity. Response to meaningful stimuli is different during sleep: for example, a person will wake up to a lower voice stimulus while hearing his own name versus hearing someone else's name [77].

The irregular sleep-wake cycle has been associated, in an ageing population, with higher cardio-vascular risk, independent of other risk factors. The metabolic consequences of frequent sleep disruption are similar to those of sleep deprivation.

Sleep latency may become problematic after frequent disruptions of the sleep and this affects the duration of sleep after the shift, acting in synergy with the short sleep duration.

Clinical evaluation: Anamnesis should focus on the duration, difficulty in sleep initiation and on the napping habits. Complaints such as diminished intellectual performance, tiredness, loss of interest, reduced level of attention or concentration on tasks, increased number of mistakes and work accidents should always raise the awareness about a possible sleep disorder. Dangerous habits, such as smoking, alcohol and auto-medication for sleeping pills might be the clinical manifestation of the sleep problems.

In clinical practice, a *structured interview* is very helpful for all employees at risk. In general, the following items are investigated: bedtime (bedtime resistance, sleep onset delay), sleep behaviour (amount, snoring, leg movements, number of waking episodes) and morning wake up (time, self-induced, sleepiness during the day). The sleepiness level, for example, can be evaluated using different scales. The extreme chronotypes are assessed with the Horne-Östberg morningness-eveningness questionnaire [78]. The major items used in this questionnaire are: time of getting up and of going to bed, level of alertness, the best interval during the day for occupational or recreational activities and the perceived difficulty to perform a certain activity at different hours. A *sleep diary* for a minimum 2 weeks is a very good tool for monitoring sleep duration, day or night distribution, weekly distribution, onset and wake up time. The non-recuperated sleep debt might be the underlying cause of the complaints.

Determination of the specific sleep pattern (periodicity of sleep duration waxes and wanes) would be ideal for pacing the shifts. The *maintenance wakefulness test* is sometimes necessary to assess the ability to stay awake in soporific conditions (quietness, dim light, comfortable temperature). It is measured using the polysomnography method under standard conditions [79] and is an objective measure of the inability of some patients to maintain normal alertness during the daytime.

The first step in any sleep disorder without organic substrates is to eliminate contextual factors for the sleep latency or delayed sleep onset. Sometimes, just social adjustments or introduction of the basic recommendations for an efficient sleep routine will be enough (relaxing activities 1/2 h before going to sleep, avoid TV, computer screen light, avoid intense physical activities 3 h before going to bed, avoid caffeine containing beverages in the second part of the day, etc.). In other cases, avoiding shift work become mandatory, at least for a certain period of time, enough for recovery.

A special issue is related to the existing diabetes or cardio-vascular disease that might be aggravated by this working schedule. In some countries, national regulations allow the occupational physician to demand the employer another kind of working time. Whenever this is possible, it seems to be the best option.

3.3. Stress

There are many *definitions of stress*. From the biological perspective, the process underlying stress is initiated by stressors and implies a stress-response (bodily reaction to stress, both psychological and physical) that also creates an emotional impact. If this impact is emotionally positive, the perception is of a positive or motivational stress; the negative emotional

impact defines the distress. In response to chronic stressor stimulation, activation of the hypothalamic-pituitary-adrenal axis and of the sympathetic nervous system increases cortisol and adrenaline systemic levels, heart rate and blood pressure. These biological effects have been used to measure, objectively, the level of stress. Chronic high levels of counter regulatory hormones (cortisol and adrenaline) are a plausible biological explanation of the insulin resistance related to stress. Indeed, in individuals with metabolic syndrome, levels of cortisol metabolites and catecholamine are found significantly higher than in controls, with a reduced cardiac frequency variability related to the higher sympathetic tonus, and high levels of inflammatory markers (protein C and IL6); the psychosocial factors are estimated to explain around 13% of these effects [80]. Negative emotions have an acute impact on the inflammatory and vasoconstrictor tonus; an interesting comparison between acute myocardial infarction precipitated by a stressful event and the ones without a distinguished negative event showed a significantly higher monocyte chemoattractant protein-1 (MCP-1) and endothelin-1 (ET-1) in the first group [81]. Distress also influences the spontaneous individual strategies to cope with stress, leading to unhealthy behaviours such as smoking, bad nutrition, low physical activity and addictions [82].

Occupational stress is considered the second most common work-related health problem, affecting 28–40% of the workforce [83, 84]. Several conceptual models have been developed to analyse the level of occupational stress. The theory of Karasek et al. based on the job demand and the control power or autonomy [85] focuses mainly on the personal control and short-term stressors. Siegrist et al. [86] created a model based on the effort-reward balance, with more emphasis on social equity and on long-term perspective [87]. A third model, the job demand-resources model, has been proposed by Bakker and Demerouti [88], challenging previous approaches for treating the diverse working conditions in a similar and general way; this model considers that even if job demands are relatively high, they will not generate distress if the organization provides the functional frame for achieving the goals, reducing by that the physical and psychological cost of the employee. Even if there are certain overlaps in these models, their focuses are different: while the Karasek et al.'s model focuses on the psychobiological impact on the threatened control, Siegrist et al. changed the accent on the threatened reward and Bakker and Demerouti have highlighted the threatened support.

In clinical research and practice, the measurement instrument of the perceived stress is the questionnaire. Depending on the conceptual model used, the items are different: for the first model, the quantity of work, the time constraints, the autonomy in making decisions, including the creativity and the possibility of skills development are the main items. The effort-reward model uses questions to assess the time constraint, the responsibility, competitiveness and fairness of treatment inside the organization, the job promotion perspectives, respect and esteem, job status, security of employment and external motivators (adequate salary). The demand-resource model claims a specific analysis by the type of activity and organization resources and hence, there is no a general questionnaire proposed. These instruments have been enriched, in certain studies, with specific items related to sleep disorders, fatigue, sufficiency of recovery time, and posttraumatic stress (the psychological injury risk indicator) [89], and they continue to evolve, creating much difficulty in the comparison of data.

Occupational stress is closely related to job dissatisfaction. Job satisfaction has a multitude of determinants, from job content, to values and leadership of the organization, management style (participative or authorship) and organizational culture (supportive or punitive, leadership role model, career paths definition, co-worker support, salary benefits, recognition, work-life balance, etc.). Nowadays, when competitiveness demands frequent improvements in the organization, proper change management processes became increasingly important in maintenance of the job satisfaction. Change is stressful if there is not enough planning, supportive leadership and if it is very frequent, creating the feeling of instability [90]. It is the management duty to find possibilities for job enrichment, job enlargement or job rotation in the process of change to increase the internal motivation and the meaningfulness of the job. But the rapid development of work and constant training create high pressure on employees and reduce family and recovery/recreational time, no matter how motivated the employee is. It can be only partially compensated, sometimes, by a perceived increase of the quality of family time through the actualization of the new skills in different social contexts.

Employees quit managers, not companies but they also leave if there is a misfit between their competencies and skills and the tasks and goals requested (poor recruitment process), if there is role ambiguity or too little decision power. Work time arrangements have several aspects in which the management of the organization can intervene to reduce stress and to increase job satisfaction and productivity such as shift work scheduling, number and flexibility of the work hours and overtime. By simply forward-rotating the shift system, the mean number of days on which the workers reported sleepiness, the heart rate at rest and the systolic blood pressure decreased significantly [91]. Longer working hours lead to increase in smoking, drinking and weight gain [92], affect the need to recovery time and create psychological distress [93]. The *Karoshi* phenomenon ('death from overwork', in Japanese, associated with more than 60 h/work month) is rarely recognized as such in western societies [94], but the silent, progressive association between insulin resistance and stress might be, interpreted, in fact, as part of the same picture.

Particularly in services, flexibility in both time and location is a new way of working that shifts the organization from standard, repetitive procedures to project-based short-term activities. In certain developed countries, such as Sweden, only 16% of jobs are regulated in terms of time, space, process flow and coordination, performance evaluation and peer interrelation, with differences between private (more flexibility) and public sectors, activity domains (services more than production) and gender: women more frequently employed in positions with higher levels of regulation than men [95]. Flexibility has certain advantages for stress reduction: it fulfils the need for autonomy and decision empowerment and gives the employee the perception of the self-determination of his working time, in terms of duration and daytime distribution. On the other hand, flexibility creates an opportunity for the employee to stay in a permanent 'switch on' work activity, predisposes to isolation, lack of supervision and peer support and reduces the development of the interpersonal skills and relations. These can lead to burnout or to unhealthy behaviours (e.g. snacking, increase consumption of sweet beverages, smoking) [96]. Family working balance is more and more important in dual-career couple societies and effort recovery time and psychological detachment from work are central in maintaining this equilibrium.

Relationship between IR and occupational stress: To answer this question, a comprehensive systematic review [97] of the prospective studies linking stress and metabolic syndrome found a positive relation. Gender seems to be important in terms of stressors: for women, the main risk factor was job strain, whereas for men the feeling of justice at work was the main protective factor. When assessing the individual components of the metabolic syndrome, weight gain was inconstantly associated with job stress, but low justice at work was associated with high triglycerides in men and job strain with high cholesterol in both sexes. Development of diabetes in men was in direct relation with higher demands. Job insecurity and low feeling of justice in men and effort-reward imbalance in women were associated with an increased high blood pressure risk. Low effort and low engagement lead to increased dissatisfaction towards job and in medium term might lead to a certain degree of depression. At the opposite end, the low effort and low engagement are overcommitment and this attitude also amplifies the adverse health effects; overcommitment is expressed by a continuous thinking of work during the normal social/relaxing time, with an impossible switch off from work. From psychological perspective, it underlies the need for approval and recognition at work [98]. The most frequent manifestations of overcommitment are burn out, psychological distress, musculoskeletal complaints and self-rated poor health [99, 100]. In a large Dutch study, overcommitment increased by more than 20 times the risk of emotional exhaustion in workers recording an imbalance between effort and reward compared to those without overcommitment [101].

Direct life-threatening jobs (policemen, firefighters, sailors) are well-known stressing conditions. Policemen has a high score in demand-control-support scale and in the effort-reward imbalance scale and these scores could explain the more than doubled OR of metabolic syndrome, particularly for the hypertriglyceridemia component [102, 103]. Working in confined spaces is another well-recognized stressor: after only 5 months of sailing around the world, in a rather small ship with only 16 persons an equipage, BMI, HbA1C, basal insulinemia and triglyceride level [104] significantly increased. In these particular working conditions, the circadian rhythm modification, the changing working environment, the rapidity in decision process, the loss of control on own private life and private activities and the anxiety in front of unknown future conditions were considered as additional stressors.

But occupational stress does not require only extreme conditions. A large British study, including more than 10,000 civil servants, using the demand control evaluation method in repetitive measurements, concluded that more than 14 years of chronic stress exposure increases the risk of metabolic syndrome by two in men and by five in women [105].

If job might be stressful, nevertheless losing job or job insecurity is another very stressful event. This association is supported by many studies [106–108]. Moreover, the relationship between the numbers of financial negative events in one's life was independently related to insulin resistance in a large, general population study in Finland [109].

Health risk factors (smoking, drinking, low physical activity, increased BMI) are more frequent in public sector employees perceiving low rewards or low occupational effort [89]. While low-reward perception is intuitively link with dissatisfaction, low effort should be expected to generate the opposite; however, in a large-scale meta-analysis, it has been found that low effort and low engagement reduce the job satisfaction and increase the level

of depression [110]. These occupational stressors generate coping behaviours for the job in satisfaction and depression. However, in both prospective and cross-sectional studies, the strength of the association could be minimized by the variety of individual reactions to stress, reducing the statistical significance for each behaviour.

Stress and nutrition relationship: As nutrition is so important in the insulin resistance physiopathology, the effect of stress on eating behaviours has been largely studied. In an ideal homeostatic balance, appetite and eating should be driven by the internal eating cues (hunger and satiety). But food is also a basic primary reward and in our food abundant modern society, numerous other external factors trigger eating. Emotional food choices are certainly influenced by the level of the stress hormones; other factors, such as gender, age and cultural modulates this behaviour. Female tends to be influenced more by the negative emotions. For example, in young women, binge eating has been related to executive function impairment such as cognitive impulsivity, insufficient planning and higher levels of cortisol release in relation to stressors. The tendency towards depression is also high among them [111]. On the contrary, men are increasing their portions, when experiencing positive emotions [112].

Stressed students and adolescents generally eat less healthy [113], and this attitude is also more influenced by negative emotions [33]; body image is the main concern [114] at this age that opposes the emotional food intake. Adults may improve emotion control and, in general, increase the health concerns about food; in this context, in healthy subjects, the positive emotions become more powerful in leading eating behaviour. Apparently, even liking for specific taste changes with age: salt liking increases and sweet taste decreases [115]. Emotional eating is frequently related to sweet taste liking that is more expressed in women [115, 116].

Stress and eating dysregulation or a poor differentiation between internal eating cues (hunger, hypoglycaemia) and external stressors are closely related and may be expressed in different personalities either as overeating or as fasting. Eating dysregulation is more frequent in persons having difficulties of identification own feelings and emotions (alexithymia) [117]. High-caloric foods (snacks) are convenient food for this type of behaviour; indeed, in experimental induced stress, people often consume more sweet fat foods than usually [118]. There is a biological explanation of this consumption, as such foods increase the serotonin level that improves mood. The mood improvement creates a positive, but medium- and long-term unhealthy, feedback loop, that contributes to permanentize the behaviour. Emotional eating has been explained by inadequate parenting practices during childhood: an increase control and restriction of certain foods for fear of weight gain (the 'dietary restraint model') or usage of foods to emotionally comfort ('affective state model') are influencing the adult emotional eating and create a significant role for the food-reward behaviour in adults [119]. Recent studies highlighted the genetic role: an increased number of central dopaminergic receptor 2 and a melatonin receptor allele are a risk factor for emotional eating and energy intake under stressful conditions [120, 121].

Even if link between stress and insulin resistance seems to be well argued by the existing data, some limitations of these studies should be highlighted. The comparability and reproducibility of data are difficult because there are a variety of instruments used to measure stress. Another limitation is that most of these studies (particularly the large ones) are studies

from the public sector, and might not be fully transposed into the private sector, where competitiveness might be higher and job employment less predictable. Even less is known about small or family enterprises that have specificities in organization, human interrelation and high interference between family and occupational stress. Personal (non-published) experience with local small-medium size firms showed that where there is tendency of preferred employment of relatives, there are additional stress factors, work inequities and general level of stress perception inside a firm.

Fortunately we still have large prospective studies confirming the risk between different components of the metabolic syndrome and stress perception. Even so, the methodology of these studies could be challenged: in many studies, there is no follow-up of the drop off (quitting the job, for example, because of the high stress, whether there is a job demand/job control induced distress or an effort-reward imbalance). If we would have the methodology to estimate in a proper way the healthy worker effect maybe the risk will be even higher, but this a common bias with all occupational risk hazards. When assessing the nutritional effect of job stress, almost all studies examined the increased in BMI and none have considered or measured the dual effect of job stress on increasing or decreasing weight. Hence, we cannot conclude whereas this would influence or not the final result.

Clinical approach: There are many types of questionnaires used to evaluate the level of stress. Selecting the most suitable instrument for an organization is sometimes difficult and involvement of psychologists and human resources specialists might be of great help. In individual assessment, the detailed anamnesis, the working time and the perceived complexity of job demands are more easily revealed during the consultation. As in other occupational medicine consultations, attention should be paid to repetitive group complaints that underline a common organizational problem. Collaboration with the line managers and human resource management in a stress reduction programme is essential.

Health status also influences the employment choice and the occupational specialist has a role in modulating this choice. For example, fatigue is one of the reasons people quit shift work and organic causes should not be omitted before claiming stress as the causative factor. Clinical examination and exchange of medical information with the general practitioner are beneficial in finding the best possible solution for the employee.

3.4. Chemical hazards and air pollution

Assessing the health impact of toxic exposure is an important part of the occupational medicine practice. During the past decades, industrial exposure to chemicals has been gradually reduced in many countries by better safety measures, but low level constant and long-term exposure are still a matter of concern for many industrial and service workplaces. Occupational exposure to heavy metals occurs in many industries not only in the metal industry (mainly from electric air furnace or welding), metal refineries, battery production, colorant and pigment industries, smelter operations, radiator repair shops, mining, construction, but also in the modern days in the recycling industry and copying machine chambers. Indoor pollution and environmental exposure (from drinking water, outdoor pollution) are other possible sources of particles and chemical exposure linked to insulin resistance.

Heavy metals: Numerous studies indicated an association between heavy metals and insulin resistance (cardio-vascular diseases and diabetes) [122–127] explained by their direct effects on ROS production and inflammatory mediated cellular and systemic damage, decreased mitochondrial function with reduction of the capacity for fatty acid oxidation, inhibition of lipoprotein lipase in adipocytes and β -cells destruction [128]. Maternal gestational diabetes is also increased by heavy metal exposure [129].

Arsenic is well known for its carcinogenic, skin and neurological action. Arsenic epidemiological data about diabetes and cardio-vascular effects is quite extensive and comes from very different countries, both developed and under developed ones [130, 131]. Industrial activity and drinking water are the major source of exposure [132]. In drinking water, levels higher than 150 mg/L are positively associated with diabetes. Lower drinking water levels gave contradictory results [133, 134] in different studies, most probably needing a longer term exposure to produce the negative health effects. Arsenic is a highly reactive metal. Inside hepatocytes, the methylation of arsenate and arsenite generates monomethylarsenate (MMA) and dimethylarsinate (DMA), the latter being less reactive and more rapidly eliminated in urine. Metabolomics techniques revealed 103 urinary and 32 plasma metabolites associated with the urinary arsenic level. Interestingly, metabolites related to energy (ATP) production were common in diabetes and non-diabetes individuals, although the energy generation is different at the cellular level, insulin is active or if there is an IR. There were also other compounds affecting the citric acid cycle or the amino acid metabolism that were specifically modified only in diabetes, supporting the influence of arsenic on the diabetes status [135]. Under moderate environmental arsenic exposure [136], high inorganic urinary As and urinary DMA excretion were associated with high diabetes incidence; higher MMA had an inverse effect. Experimental data also supports an action of DMA on adipocytes (reduction of the insulin signal) and on beta cells apoptosis. Exposure burden, the type of arsenic compounds but also polymorphisms of arsenic (III) methyltransferase gene (*AS3MT*) [137] are relevant for the relation between MMA and DMA and nutrition, particularly on folate intake.

In animal experiments, arsenic has many pathophysiological effects related to insulin resistance: it reduces the intracellular insulin signal by suppressing the expression and phosphorylation of protein kinase B (PKB/Akt) and by direct inhibition of the Akt pathway [138]. Arsenic affects hepatic glucose output, increasing gluconeogenesis and up-regulates genes involved in oxidative and inflammation processes. As other heavy metals, arsenic causes mitochondrial toxicity reflected by the decreased level of the mitochondrial deacetylase, of FOXO 3a, ionic transport and membrane potential, decreased activity of manganese superoxide dismutase and increased reactive oxygen species [139]. Arsenic activates stress sensitive signalling pathways initiating pancreatic cell apoptosis [140] and oxidative stress in liver cells [141]. Arsenic has also a pro-thrombotic action and acceleration of atherosclerosis explaining the cardio-vascular effects [142, 143].

Mercury (Hg): Cross-sectional and prospective studies showing increases prevalence of diabetes [144, 145] in people exposed to Hg, which are not confirmed by others [146]. Hg is a very reactive metal: it interacts with the -SH groups of different enzymes, antioxidants

(glutathione), and even with amino acids. Substantial experimental data concluded that Hg has both direct pro-oxidant effects and a capacity to reduce the antioxidant defence mechanism [147]. These characteristics have been confirmed in high occupational exposure [148, 149]. Hg reduces glutathione, superoxide dismutase, catalase and thioredoxin reductase [150]. Denaturation of proteins increases the endoplasmic reticulum stress, initiating apoptosis. The inflammatory status is materialized by the increased levels of IL4, IL6, IL7 and TNF α and IFN γ [150]. Apart from these systemic effects, Hg has also more specific effects related to IR development: Hg has a direct toxic, pro-apoptotic effect on β -cells [151]; it decreases the insulin effects [152] and induces apoptosis in adipocytes.

Other heavy metals have also been involved in metabolic inflammation. During the cellular immune response, it was demonstrated that radicals induced by lead (Pb) disrupt the transcription signalling pathway mediated by the mitogen-activated protein kinase, nickel (Ni) induces the NLRP3-ASC-caspase 1 and As induces tyrosine kinase Src. The nuclear factor kB is activated by Pb, Ni or Hg with an increase in gene transcription for early inflammatory cytokines, such as Interleukin 1, interleukin 6 and tumour necrosis factor. Some metals, such as cadmium (Cd), can activate an inflammatory response through tissue damage induction mediated by free radicals, which results in leukocyte recruitment and cytokine secretions. Inflammation generated by metals can be reduced by metallothionein that has the ability to scavenge free radicals and bind toxic metals through the release of Zn and oxidation of SH groups [153].

From my previous personal clinical experience, in the Occupational Medicine Clinic of the Colentina University Hospital in Bucharest, I can confirm that workers from a lead battery factory, currently closed, admitted with saturnine colic, that had heavy Pb exposure, presented almost constantly a reversible hypertension syndrome.

Heavy metals and smoking: Smoking habit is one of the most important sources of cadmium in the general population. In fact, reduction of cadmium levels in the general population has been attributed to the efficiency of the anti-tobacco campaigns and policies [154]. Lead, nickel and chromium are also found in the tobacco smoke at significant levels. The relationship between smoking and diabetes is grounded on large epidemiological studies [155, 156]. Mechanisms such as impaired peripheral glucose uptake, oxidative stress, metabolic inflammation [157], and retention of the inhaled high concentration of metals (As, Al, Cd, Ni, Hg and Pb) [158, 159] have been described. A synergic effect of occupational exposure and smoking was found in smelters [160] or mining workers [159]; the common proposed pathological mechanism is the reduction of the anti-oxidative defence [160] or the increased retention of the heavy metals.

Other toxicants: Epidemiological studies brought the attention on other environmental toxicants such as persistent organic pollutants (POP), endocrine disruptors, or substances involved in the toxic oil syndrome (presumably aniline). POP is a generic term for a variety of chloride or bromide compounds, including pesticides and dioxin. Longitudinal studies showed that persons with high basal level of blood POP are at increased risk for diabetes. Studies that have also monitored the PCB were able to exclude the non-reverse effect of

diabetes [161]. The stronger evidence of these associations came from occupational exposure studies: high accidental environment contamination [162] increased the relative risk of diabetes and of cardiovascular disease. Non-accidental occupational exposure in cohorts from 12 different countries also confirmed the relationship [163]. Bisphenol A is also a POP, well known for its endocrine disruptor effects; besides other negative effects, the rate of gestational diabetes is increased in women with a higher blood level [164]. Toxic oil syndrome-registered patients have threefold increase in prevalence of diabetes, and higher incidence of metabolic syndrome [165, 166].

The intimate mechanism of these associations is not completely found and most probably has certain specificities according to the particular substance. Many pathological pathways have been implicated: the increased accumulation of abdominal fat [167], inflammatory markers [168], adipocyte cytokines, TNF α and nuclear transcription factor kappa B (NF κ B) [47], uric acid [169]. As a direct effect on the insulin signal, in animal experiments, perfluorooctane sulfonate inhibits the insulin Akt, signal leading to insulin resistance [170].

Particulate exposure: Air pollution, particularly fine and ultrafine particles, has been related to the acute incidence of cardio-vascular events and asthma attacks from inflammatory, oxidative mechanisms and pro-thrombotic mechanisms activation [171]. This correlation remained consistent after exclusion of possible confounders (education level, work exposure, alcohol, vegetable and fruit consumption, smoking or physical activity) [172]. Initially described for cardiovascular effects [171, 173], the particulate exposure includes now also the increased risk for diabetes [174, 175] and for non-alcoholic steatohepatitis [176]. Particulate matter (PM) is generally a subject of environmental health, and exposure during the day is very variable depending the microenvironment (home, kitchen, policy of smoking indoor, traffic, general level of city pollution control, occupation) [177]. However, there are also many workplaces where PM₁₀ (the respirable component), PM_{2.5} (fine particles) and the PM_{0.1} (ultrafine particles) have still high values. Road transport, energy production and distribution, waste as well as domestic exposure (cooking) are major sources in Europe [178]. Photocopiers centres, sick buildings might also have PM₁₀ and PM_{2.5} above than permissible levels.

Large cohort studies in North America showed that a 10 g/mm³ increase in PM_{2.5} was associated with 13% increases in hypertension and with 18% increase in diabetes-related mortality, concluding that current PM 2.5 US standards are too high for the long-term exposure and should be reduced [179].

In different countries or regions, the proportion and the composition of air pollution differ; in a China study, fine particles bound polycyclic aromatic hydrocarbons (PAH) and this characteristic might have a cumulative effect on IR, as it has independently associated with this risk [180]. PAH is an environmental pollutant but exposure is increased in occupations that involve bitumen and asphalt manipulation, in foundries, distilleries, painting industry and other manufactories [181, 182]. Firefighters are also exposed to significant levels. Smoking is another important PAH source and, therefore, efforts to reduce this habit in workers occupationally exposed are highly recommended.

The particulate matters IR pathophysiologic mechanisms are rather similar to the ones involved in heavy metal exposure: obesity risk increase [183], enhancement of inflammation, with higher plasma levels IL8, ICAM-1 or eosinophilic cationic protein [184], endoplasmic reticulum stress and insulin signalling interference in a dose-response manner [172]. Reduction in antioxidant defence is supported by both experimental on cell lines and from the increased effect on persons with GSTM1 null alleles [184]. Disruption of metabolic genes has also been reported [185]. Several studies proposed a link between genetic variants of ABO locus; particulate exposure and IR risk [186]. When occupational exposure to particles (particularly to asbestos, welding fumes, respiratory irritants) was added to the epidemiological model, the risk of group O subjects was enhanced [187]. There is no conclusive explanation about this association, but ABO antigens are expressed also on the von Willebrand factor, presumably increasing the prothrombotic and chronic inflammation risk [188] and the ABO locus is related to the E-selectin, ICAM-1 and TNF- α blood level [186].

Cardio-vascular and ECG modifications: Controlled particulate exposure showed high systolic [189] (or high systolic and diastolic increase [172] according to the type of particle), decreased of blood plasminogen and thrombomodulin, and an increase in the inflammatory markers: CRP and serum amyloid [190]. Diesel exhaust double-blind experiment mean peak effect on systolic blood pressure increases between 30 and 60 min. after exposure and remained high 24 h after, independent of metabolic syndrome previously existing components. Interestingly, in this experiment, the increase in blood pressure was not related to perception of exposure [189]. Persons with metabolic syndrome exposed to ultrafine particles and GSTM1 null allele are particularly at risk of altered repolarization and increased QT duration [191].

Clinical approach: Indoor air pollution measurement is part of the occupational safety procedures and whenever data are available there should be transmitted to the occupational physician. The detailed occupational anamnesis, symptoms, clinical examination and existing exposure indicators orient the medical conduct.

For heavy metals exposure, biological monitoring is recommended in workplaces or occupations at risk. The biological monitoring is included in the national recommendations for occupational medicine practice in many countries. The limit values of the exposure indicators are defined for chronic poisoning. On what concerns the metabolic risk, there is no threshold limit established yet and general population data suggest that lower limits than those for chronic poisoning would be a reasonable approach. We cannot conclude this topic, until well-designed studies to address this specific objective are conducted. On the other hand, until these studies are available and taken into consideration the increasing incidence of insulin resistance worldwide, efforts to reduce this non-classical risk factor should not be postponed.

A special concern is related to the maternal and foetal risk and to the gestational diabetes. Ideally, pregnant women should not be exposed to chemicals; if environmental contamination is not easy to avoid, at least the occupational one should be limited. Besides the negative effects of chemicals on the foetus development, the increased risk of gestational diabetes adds unfavourable short- and long-term consequences.

4. Workplace interventions

As insulin resistance implication is better understood, and its prevalence increases worldwide, the number of interventions that have been published in last decade has also increased. Most of them, even if they directly target insulin resistance are better known as workplace wellness programmes. In countries where private insurance is paid by the employer, they could be a mean to reduce the insurance plans; in others, where work-promotion activities are more and more valued by the companies, they are conducted either as a channel to encourage individuals to take preventive measures through education and screening or as employment-based activity to promote health behaviours and disease management, such as stress management, adjustable work station to reduce continuous sitting, team support groups and cafeteria/canteen provision of healthy food initiatives. Although very diverse, there are however some general characteristics of these interventions:

1. *Assessment of the health risk according to the working conditions.* Risk assessment is always the first step as the intervention has to be meaningful for the organization and should reflect the specific issues of the employees. Working conditions and health-related habits should be analysed, as multi-component interventions have given better results [160]. Sometimes, just screening programmes initiated inside the company increase awareness and initiate individual actions [161]. In works with high aerobic workload, a low cardiovascular fitness is a physical risk factor and a stressor. Increasing the aerobic fitness by initiating an exercise training programme was able to decrease the basal heart rate, the BMI and the inflammatory markers (high sensitivity protein C), to increase work efficiency and to reduce stress related to work overload. Sedentary behaviour has benefited from different ergonomically interventions [162, 163]. Some studies have emphasized the benefit of sitting interruption during the working day; efficiency of short 1–2 min every half hour of sitting break was better in terms of metabolic syndrome criteria compared to the two 15 min breaks per workday [164]. In certain circumstances, cumulative risk factors are encountered: for example, seafarers are recognized as a profession with high incidence of metabolic syndrome. An intervention including healthy cooking training for ship cooks together with improvement of the ships fitness facilities and education (anti-smoking, individual exercise, extra health checks) led to a decrease in sugar intake, increased physical fitness scores and a significantly decrease in prevalence of the metabolic syndrome [165, 166].
2. *Obtaining the maximum support of the management team.* These interventions are changing behaviour interventions; they imply not only financial cost but also time allocation, organizational changes, etc. If management is not convinced and does not actively supports the programme, making it part of the organizational culture, the chance of achieving the proposed results is considerably reduced. Presenteeism, defined as being present at work, but with a limited performance in some aspects of job related a health problem, is often a high hidden cost for employers. Improving the general health status and well-being of their employees could be, therefore, perceived as a valuable investment. Stress management programmes, in particular, need specific organizational behaviour assessment and

a proactive participation of the line and human resource management. Of course, there are some universally interventions for mental health [167], but they should be aligned with other managerial and organizational initiatives [168].

3. *Proper preparation and communication inside the company:* Ideally, some representatives of the beneficiaries should be involved in preparation of the intervention, for presenting their needs and for choosing from different possibilities of implementation the best suited for them. Communication of the aim, the means and the expected results is necessary before, during and after the intervention. It is reasonable to consider that employees are in different stages of motivation for change. Therefore, stage-based intervention for non-intenders, intenders and actors (people who have already taken action towards changing behaviour) could be more effective [169, 170]. Barriers to adherence and completion of the programme are better solved in a participatory approach [171].
4. *Adapted tools:* There is no one solution for all organizations. For some, the online support is possible and is part of their working culture; in other organizations, group or individual discussions are more effective. Computer-based interventions have a better reach, particularly in larger organizations. A Cochrane systematic review on interactive computer-based interventions for weight loss or weight maintenance in overweight or obese people found that such interventions significantly reduced body weight [172]. Combined systems with face-to-face and computer/telephone interactions are also used.
5. *Team work,* from the first steps of design until results evaluation. Nobody can expect that a transformational change can be proposed and delivered solely by the occupational physician, but it is desirable that he is an active member of the team [173]; psychologists, ergonomists, nutritionists, exercise trainers, IT and communication specialists, etc. are needed, according to the scope of the intervention. Concerning risk reduction of the metabolic risk, nutrition is a traditional risk factor for insulin resistance and trained dieticians should be involved. Nutrition knowledge make people less likely to be hypertensive compared to one with low level of healthy nutrition [174] and when combined with a physical activity programmes [175], and facilitated access to healthy food, become more effective [176]. Workplaces are excellent settings for health promotion programmes, joining all these conditions together, if there is managerial commitment and support.

5. Conclusions

The minimum requirements for a health issue to become an occupational medicine subject are a proven relation with the working conditions, and a benefit from a working place intervention. This chapter had provided arguments that insulin resistance has both characteristics. It has, however, a specificity, that in some countries had brought benefits for including it in the occupational services and in others have been an obstacle: insulin resistance shares both elements of the classical health protection and of the health promotion activities.

In terms of health protection, insulin resistance is related to medium or even low exposure to air pollution (particles, solvents) or heavy metals. The biological monitoring reference for insulin resistance is not yet defined, making difficulties in interpreting field data. Using the general population assessments might be a solution, but there is clearly a need for well-designed research in this area to reach a valid conclusion. Insulin resistance is also related to other work hazards: work schedules, sedentary behaviour and stress. Concerning the work schedules, we benefit of many data, experience and general recommendations about working on shifts; flexible working time is a new challenge and probably needs more individual solutions. Occupational stress is recognized as one of the most important risk factor of the modern society and, even if it is more in the occupational psychology field, it has to be integrated in the occupational medicine service, as many other medical conditions, except insulin resistance, are aggravated by stress.

In terms of health promotion, prevention of insulin resistance and its major consequences (the cardio-vascular disease and type-2 diabetes) are perceived as part of the healthy lifestyle and well-being programmes. Healthy nutrition, smoking cessation, physical activity and sleep hygiene are the pillars of these programmes. The appropriate and culturally adapted design increases the adherence to these programmes; their real success is the integration of the healthy behaviours and the reduction of obesity and insulin resistance-related mortality. Management support is needed in the preliminary phases (approving and promoting the programme) but also during the implementation phase, as many of these activities request working condition adaptations. Better employee engagement is obtained by involving them from the early steps of the programme.

Therefore, assessment of the risk, screening and workplace intervention programmes to reduce insulin resistance incidence are part of the occupational medicine activity and should be included in the occupational health service provision. Further research to develop tools for on appropriate individual and organizational assessment and strategies for interventions is needed.

Author details

Marina Oțelea

Address all correspondence to: dr.marinaotelea@gmail.com

Clinical Department 2, University of Medicine and Pharmacy “Carol Davila”, Bucharest, Romania

References

- [1] WHO. Media centre. Cardio-vascular diseases [Internet]. Available from: <http://www.who.int/mediacentre/factsheets/fs317/en/> [Accessed 15.05.2016].

- [2] WHO Media centre. Diabetes [Internet]. Available from: <http://www.who.int/mediacentre/factsheets/fs312/en/> [Accessed 15.05.2016].
- [3] Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: How education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Pub Health*. 1992;**82**:816–820.
- [4] Santos A, Ebrahim S, Barros H. Gender, socio-economic status and metabolic syndrome in middle-aged and old adults. *BMC Public Health*. 2008;**8**:62. DOI: 10.1186/1471-2458-8-62.
- [5] Al-Daghri NM, Alkharfy KM, Al-Attas OS, Khan N, Alfawaz HA, Alghanim SA, et al. Gender-dependent associations between socioeconomic status and metabolic syndrome: A cross-sectional study in the adult Saudi population. *BMC Cardiovasc Disord*. 2014;**14**:51. DOI: 10.1186/1471-2261-14-51.
- [6] Langenberg C, Kuh D, Wadsworth MEJ, Brunner E, Hardy R. Social circumstances and education: Life course origins of social inequalities in metabolic risk in a prospective national birth cohort. *Am J Pub Health*. 2006;**96**:2216–2221. doi:10.2105/AJPH.2004.049429.
- [7] Schooling CM, Jiang CQ, Lam TH, Zhang WS, Cheng KK, Leung G. M. Life-course origins of social inequalities in metabolic risk in the population of a developing country. *Am J Epidemiol*. 2008;**167**:419–428.
- [8] Gupta R, Deedwania PC, Sharma K, Gupta A, Guptha S, Achari V et al. Association of educational, occupational and socioeconomic status with cardiovascular risk factors in Asian Indians: A cross-sectional study. Gupta V. *PLoS ONE*. 2012;**7**:e44098. DOI:10.1371/journal.pone.0044098.
- [9] Lucove JC, Kaufman JS, James SA. Association between adult and childhood socioeconomic status and prevalence of the metabolic syndrome in African Americans: The pitt county study. *Am J Public Health*. 2007;**97**:234–236. DOI:10.2105/AJPH.2006.087429.
- [10] Eibner C, Evans WN. Relative deprivation, poor health habits, and mortality. *J Hum Res*, 2005;**40**:591–620.
- [11] Popkin BM, Gordon-Larse P. The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord*. 2004;**28**:S2–S9. DOI:10.1038/sj.ijo.0802804.
- [12] Omran AR. The epidemiologic transition: A theory of the epidemiology of population change. *The Milbank Q*. 2005;**83**:731–757. DOI:10.1111/j.1468-0009.2005.00398.x.
- [13] Cornman JC, Gleib DA, Goldman N, Ryff CD, Weinstein M. Socioeconomic status and biological markers of health: An examination of adults in the United States and Taiwan. *J Aging Health*. 2015;**27**:75–102.
- [14] Dallongeville J, Cottel D, Ferrières J, Arveiler D, Bingham B, Ruidavets JB, et al. Household income is associated with the risk of metabolic syndrome in a sex-specific manner. *Diabetes Care*. 2005;**28**:409–415. DOI: 10.2337/131224.

- [15] Davila EP, Florez H, Fleming LE, et al. Prevalence of the metabolic syndrome among U.S. workers. *Diabetes Care*. 2010;**33**:2390–2395. DOI:10.2337/dc10-0681.
- [16] Alegría E, Cordero A, Laclaustra M, Grima A, León M, Casanovas J. Prevalence of metabolic syndrome in the Spanish working population: MESYAS registry. *Rev Esp Cardiol*. 2005;**58**:797–806.
- [17] Lee W, Yeom H, Yoon JH, Jung PK, Lee JH, Seok H, et al. Metabolic outcomes of workers according to the International Standard Classification of Occupations in Korea. *Am J Ind Med*. 2016;**59**:685–694.
- [18] Church TS, Thomas DM, Tudor-Locke C, Katzmarzyk PT, Earnest CP, Rodarte RQ. Trends over 5 decades in U.S. occupation-related physical activity and their associations with obesity. *PLoS ONE*. 2011;**6**:e19657.
- [19] Bertrais S, Beyeme-Ondoua, JP, Czernichow S, Galan P, Hercberg S, Oppert JM. Sedentary behaviors, physical activity, and metabolic syndrome in middle-aged French subjects. *Obes Res*, 2005;**13**:936–944. DOI:10.1038/oby.2005.108.
- [20] Herman MA, Kahn BB. Glucose transport and sensing in the maintenance of glucose homeostasis and metabolic harmony. *JCI*. 2006;**116**(7):1767–1775.
- [21] Poehlman ET, Dvorak RV, DeNino WF, Brochu M, Ades PA. Effects of resistance training and endurance training on insulin sensitivity in nonobese, young women: A controlled randomized trial. *J Clin Endocrinol Metab*. 2000;**85**:2463–2468. <http://doi.org/10.1210/jcem.85.7.6692>.
- [22] Menshikova EV, Ritov VB, Fairfull L, Ferrell RE, Kelley DE, Goodpaster BH. Effects of exercise on mitochondrial content and function in aging human skeletal muscle. *J Gerontol A, Biol Sci Med Sci*. 2006;**61**:534–540.
- [23] Balakrishnan VS, Rao M, Menon V, et al. Resistance training increases muscle mitochondrial biogenesis in patients with chronic kidney disease. *Clin J Am Soc Nephrol*. 2010;**5**:996–1002.
- [24] Melov S, Tarnopolsky MA, Beckman K, Felkey K, Hubbard A. Resistance exercise reverses aging in human skeletal muscle. *PLoS One*. 2007;**2**:e465.
- [25] Giresi P, Stevenson E, Theilhaber J, Koncarevic A, Parkington J, Fielding R, et al. Identification of a molecular signature of sarcopenia. *Physiol Genomics* 2005;**21**:253–263.
- [26] Yin H, Price F, Rudnicki MA. Satellite cells and the muscle stem cell niche. *Physiol Rev*. 2013;**93**:23–67. DOI:10.1152/physrev.00043.2011.
- [27] Civitarese AE, Carling S, Heilbronn LK, Hulver MH, Ukropcova B, Deutsch WA. Calorie restriction increases muscle mitochondrial biogenesis in healthy humans. *PLoS Medicine*. 2007;**4**:485–494.

- [28] Laaksonen DE, Lakka HM, Lynch J, Lakka TA, Niskanen L, Rauramaa R, et al. Cardiorespiratory fitness and vigorous leisure-time physical activity modify the association of small size at birth with the metabolic syndrome. *Diabetes Care*. 2003;**26**:2156–2164.
- [29] Randle PJ, Garland PB, Hales CN, Newsholme EA. The glucose fatty-acid cycle. Its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet*. 1963;**1**:785–789.
- [30] Wells J, Siervo M. Obesity and energy balance: Is the tail wagging the dog? *Eur J Clin Nutr*. 2011;**65**:1173–1189. DOI:10.1038/ejcn.2011.132.
- [31] Cunningham JJ. Body composition as a determinant of energy expenditure: A synthetic review and a proposed general prediction equation. *Am J Clin Nutr*. 1991;**54**:963–969.
- [32] Johnstone AM, Murison SD, Duncan JS, Rance KA, Speakman JR. Factors influencing variation in basal metabolic rate include fat-free mass, fat mass, age, and circulating thyroxine but not sex, circulating leptin, or triiodothyronine. *Am J Clin Nutr*. 2005;**82**:941–948.
- [33] Dvorak RV, DeNino WF, Ades PA, Poehlman ET. Phenotypic characteristics associated with insulin resistance in metabolically obese but normal weight young women. *Diabetes*. 1999;**48**:2210–2214.
- [34] Bays HE. Adiposopathy. Is “sick fat” a cardiovascular disease? *J Am Coll Cardiol*. 2011;**57**:2461–2473.
- [35] Sisson SB, Camhi SM, Church TS, Martin CK, Tudor-Locke C, Bouchard C, et al. Leisure time sedentary behavior, occupational/domestic physical activity, and metabolic syndrome in U.S. men and women. *Metab Syndr Relat Disord*. 2009;**7**:529–536.
- [36] Pietroiusti A, Magrini A, Coppeta L, Barone S, Sed B, Manuli G, et al. Metabolic syndrome among operators using video display units in call centers. *Scand J Work, Environ Health*. 2007;**Suppl 33**(3):49–53.
- [37] Shin SY, Lee CG, Song HS, Kim SH, Lee HS, Jung MS, et al. Cardiovascular disease risk of bus drivers in a city of Korea. *Ann Occup Environ Med*. 2013;**25**:34.
- [38] Leischik R, Foshag P, Strauss M, Littwitz H, Garg P, Dworak B, et al. Aerobic capacity, physical activity and metabolic risk factors in firefighters compared with police officers and sedentary clerks. *PLoS ONE*. 2015;**10**: e0133113. DOI:10.1371/journal.pone.0133113.
- [39] Parry S, Straker L. The contribution of office work to sedentary behaviour associated risk. *BMC Public Health*. 2013;**13**:1–10. <http://doi.org/10.1186/1471-2458-13-296>.
- [40] Healy G, Dunstan DW, Salmon J, Cerin E, Shaw J, Zimmet P, et al. Breaks in sedentary time. Beneficial associations with metabolic risk. *Diabetes Care*. 2008;**31**:661–666.
- [41] Bey L, Hamilton MT. Suppression of skeletal muscle lipoprotein lipase activity during physical inactivity: A molecular reason to maintain daily low-intensity activity. *J Physiol*. 2003;**551**:673–682.

- [42] Hamilton M, Hamilton D, Zderic T. Role of Low Energy Expenditure and Sitting in Obesity, Metabolic Syndrome, Type 2 Diabetes, and Cardiovascular Disease. *Diabetes*. 2007;**56**:2655–2667.
- [43] Mabry RM, Winkler EH, Reeves MM, Eakin EG, Owen N. Associations of physical activity and sitting time with the metabolic syndrome among Omani adults. *Obesity*. 2012;**20**:2290–2295.
- [44] Stiller JW, Postolache TT. Sleep-wake and other biological rhythms: Functional neuroanatomy. *Clin Sports Med*. 2005;**24**:205–235.
- [45] Wong SN, Halaki M, Chow CM. The periodicity of sleep duration — an infradian rhythm in spontaneous living. *Nat Sci Sleep*. 2013;**5**:1–6.
- [46] Toh KL. Basic science review on circadian rhythm biology and circadian sleep disorders. *Ann Acad Med Singapore*. 2008;**37**:662–668.
- [47] Nakagawa H, Sack RL, Lewy AJ. Sleep propensity free-runs with the temperature, melatonin, and cortisol rhythms in a totally blind person. *Sleep*. 1992;**15**:330–336.
- [48] Reiter RJ, Tan DX, Mayo JC, Sainz RM, Leon J, Czarnecki Z. Melatonin as an antioxidant: Biochemical mechanisms and pathophysiological implications. *Acta Biochim Pol*. 2003;**50**:1129–1146.
- [49] Hense S, Barba G, Pohlabein H, De henauw S, Marild S, Molnar D et al. Factors that influence sleep duration in European children. *Sleep*. 2011;**34**(5):633–639.
- [50] Leng Y, Wainwright NWJ, Cappuccio FP, Surtees PG, Luben R, Wareham Net al. Self-reported sleep patterns in a British population cohort. *Sleep Med*. 2014;**15**:295–302. DOI:10.1016/j.sleep.2013.10.015.
- [51] Hao YL, Zhang B, Jia FJ, Li XL, Tang Y, Ren YZ et al. A three-phase epidemiological study of short and long sleepers in a middle-aged Chinese population: Prevalence and characteristics. *Braz J Med Biol Res*. 2014;**47**:157–165.
- [52] Touchette E, Côté SM, Petit D, Liu X, Boivin M, Falissard B, et al. Short nighttime sleep-duration and hyperactivity trajectories in early childhood. *Pediatrics*. 2009;**124**:e985.
- [53] He Y, Jones CR, Fujiki N, Xu Y, Guo B, Holder JL Jr, et al. The transcriptional repressor DEC2 regulates sleep length in mammals. *Science (NY)*. 2009;**325**:866–870. DOI:10.1126/science.1174443.
- [54] Lakerveld J, Mackenbach JD, Horvath E, Rutters F, Compernelle S, Bárdos H, et al. The relation between sleep duration and sedentary behaviours in European adults. *Obes Rev*. 2016;**17**:62–67.
- [55] Archer SN, Laing EE, Möller-Levet CS, van der Veen DR, Bucca G, Lazar AS, et al. Mistimed sleep disrupts circadian regulation of the human transcriptome. *Proc Natl Acad Sci USA*. 2014;**111**:E682–E691.

- [56] Partinen M, Kaprio J, Koskenvuo M, Putkonen P, Langinvainio H. Genetic and environmental determination of human sleep. *Sleep*. 1983;**6**:179–185.
- [57] Maurovich-Horvat E, Pollmächer T, Šonka K. The effects of sleep and sleep deprivation on metabolic, endocrine and immune parameters. *Prague Med Rep*. 2008;**109**:275–285.
- [58] Peschke E, Frese T, Chankiewitz E, Peschke D, Preis U, Schneyer U, et al. Diabetic Goto Kakizaki rats as well as type 2 diabetic patients show a decreased diurnal serum melatonin level and an increased pancreatic melatonin-receptor status. *J Pineal Res*. 2006;**40**:135–143.
- [59] Frellick F. Melatonin links gene variant to Type 2 diabetes. 2016. Available from: <http://www.medscape.com/viewarticle/863531> [Accessed 2016-08-25]
- [60] Engle-Friedman M. The effects of sleep loss on capacity and effort. *Sleep Science*. 2014;**7**:213–224. <http://doi.org/10.1016/j.slsci.2014.11.001>.
- [61] Schmid SM, Hallschmid M, Jauch-Chara K, Bandorf N, Born J, Schultes B. Sleep loss alters basal metabolic hormone secretion and modulates the dynamic counterregulatory response to hypoglycemia. *J Clin Endocrinol Metab*. 2007;**92**:3044–3051. <http://doi.org/10.1210/jc.2006-2788>
- [62] Broussard JL, Ehrmann DA, Van Cauter E, Tasali E, Brady MJ. Impaired insulin signaling in human adipocytes after experimental sleep restriction: A randomized, crossover study. *Ann Intern Med*. 2012;**157**:549–557.
- [63] Mander BA, Colechia E, Spiegel KS, Van Cauter E. Short Sleep: A risk factor for insulin resistance and obesity. *Sleep*. 2001;**Suppl 24**:A74–75.
- [64] Leproult R, Deliens G, Gilson M, Peigneux P. Beneficial impact of sleep extension on fasting insulin sensitivity in adults with habitual sleep restriction. *Sleep*. 2015;**38**:707–715. DOI:10.5665/sleep.4660.
- [65] Hampton SM, Morgan LM, Lawrence N, et al. Postprandial hormone and metabolic responses in simulated shift work. *J Endocrinol*. 1996;**151**:259–267.
- [66] Spiegel K, Tasali E, Penev P, van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels and increased hunger and appetite. *Ann Intern Med*, 2004;**141**:846–850.
- [67] Canuto R, Pattussi MP, Macagnan JBA, Henn RL, Olinto MTA. Metabolic syndrome in fixed-shift workers. *Rev Saúde Pública*. 2015;**49**. <http://doi.org/10.1590/S0034-8910.2015049005524>.
- [68] Scheer FA, Hilton MF, Mantzoros CS, Shea SA. Adverse metabolic and cardiovascular consequences of circadian misalignment. *PNAS*. 2009;**106**:4453–4458.
- [69] Litinski M, Scheer FA, Shea SA. Influence of the circadian system on disease severity. *Sleep Med Clin*. 2009;**4**:14v3–163. DOI:10.1016/j.jsmc.2009.02.005.
- [70] Garaulet M, Ordovás JM, Madrid JA. The chronobiology, etiology and pathophysiology of obesity. *Int J Obes*. 2010;**34**:1667–1683.

- [71] Schoeller D. Entrainment of the diurnal rhythm of plasma leptin to meal timing. *J Clin Invest.* 1997;**100**:1882–1887.
- [72] Marcheva B, Ramsey KM, Buhr ED, Kobayashi Y, Su H, Ko CH et al. Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes. *Nature* 2010;**466**:627–663.
- [73] Maury E, Ramsey M, Bass KJ. Circadian rhythms and metabolic syndrome: From experimental genetics to human disease. *Circ Res.* 2010;**106**(3):447–462.
- [74] Lehman JJ, Brager PM, Kovacs A, Saffitz JE, Medeiros DM, Kelly DP. PPAR γ coactivator-1 (PGC-1) promotes cardiac mitochondrial biogenesis. *J Clin. Invest.* 2000;**106**:847–856.
- [75] Taheri S, Lin L, Austin D, Young T, Mignot E. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Medicine.* 2004;**1**:e62.
- [76] Imaki M, Hatanaka Y, Ogawa Y, Yoshida Y, Tanada S. An epidemiological study on relationship between the hours of sleep and life style factors in Japanese factory workers. *J Physiol Anthropol Appl Human Sci.* 2002;**21**:115–120.
- [77] Oswald I, Taylor AM, Treisman M. Discriminative responses to stimulation during human sleep. *Brain.* 1960;**83**:440–453.
- [78] Horne J, Östberg O. A self-assessment questionnaire to determine morningness eveningness in human circadian rhythms. *Int J Chronobiol.* 1976;**4**:97–100.
- [79] Doghramji K, Mitler MM, Sangal R, Shapiro C, Taylor S, Walsleben J, et al. A normative study of the maintenance of wakefulness test (MWT). *Electroencephalogr Clin Neurophysiol.* 1997;**103**:554–562.
- [80] Brunner EJ, Hemingway H, Walker BR, Page M, Clarke P, Juneja M, et al. Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome, nested case-control study. *Circulation.* 2002;**106**:2659–2665.
- [81] Wilbert-Lampen U, Nickel T, Leisner D, Guthlin D, Matis T, Volker C, et al. Modified serum profiles of inflammatory and vasoconstrictive factors in patients with emotional stress-induced acute coronary syndrome during world cup soccer 2006. *J Am Coll Cardiol.* 2010;**55**:637–642.
- [82] Kouvonen A, Kivimäki M, Virtanen M, Heponiemi T, Elovainio M, Pentti J, et al. Effort-reward imbalance at work and the co-occurrence of lifestyle risk factors: Cross-sectional survey in a sample of 36,127 public sector employees. *BMC Public Health.* 2006;**6**:24. DOI:10.1186/1471-2458-6-24.
- [83] European agency for safety and health at work. Facts 22. Work related stress. 2002. Available from: <https://osha.europa.eu/en/tools-and-publications/publications/factsheets/22> [Accessed: 2016-05-15]
- [84] Sauter S, Murphy L, Colligan M, Swanson N, Hurrell J, Scharf F, et al. Stress at work. Available from: <http://www.cdc.gov/niosh/docs/99-101/>. [Accessed: 2016-05-15]

- [85] Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick, B. The job content questionnaire (JCQ): An instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol.* 1998;**3**:322–355.
- [86] Siegrist J, Starke D, Chandola T, Godin I, Marmot M, Niedhammer I, et al. The measurement of effort-reward imbalance at work: European comparisons. *Soc Sci Med.* 2004;**58**:1483–1499.
- [87] Pelfrene E, Clays E, Moreau M, Mak R, Vlerick P, Kornitzer M, De Backer G. The job content questionnaire: Methodological considerations and challenges for future research by. *Arch Public Health.* 2003;**61**:53–74.
- [88] Bakker AB, Demerouti E. The job demands resources model: State of the art. *journal of managerial psychology.* 2007;**22**:309–328. <http://doi.org/10.1108/02683940710733115>.
- [89] Magnavita N. Work-related psychological injury is associated with metabolic syndrome components in apparently healthy workers. *Plos One.* 2015;**10**:e0130944.
- [90] Rafferty AE, Griffin MA. Perceptions of organizational change: A stress and coping perspective. *J Appl Psychol.* 2006;**91**:1154–1162.
- [91] Viitasalo K, Kuosma E, Laitinen J, Härmä M. Effects of shift rotation and the flexibility of a shift system on daytime alertness and cardiovascular risk factors. *Scand J Work, Environ Health.* 2008;**34**:198–205. <http://doi.org/10.5271/sjweh.1228>.
- [92] Shields M. Long working hours and health. *Health Rep.* 1999;**11**:33–48.
- [93] De Raeye L, Jansen NWH, Kant IJ. Health effects of transitions in work schedule, work hours and overtime in a prospective cohort study. *Scand J Work, Environ Health.* 2007;**33**:105–113.
- [94] Nishiyama K, Johnson JV. Karoshi--death from overwork: Occupational health consequences of Japanese production management. *Int J Helath Serv.* 1997;**27**:625–641.
- [95] Alvin M, Mellner C, Movitz F, Aronsson G. The diffusion of flexibility: Estimating the incidence of low-regulated working conditions. *Nordic Journal of Working Life Studies.* 2013;**3**:99–116.
- [96] Janssen D, Nachreiner F. Health and psychosocial effects of flexible working hours. *Rev Saude Publica.* 2004;**Suppl 38**. <http://dx.doi.org/10.1590/S0034-89102004000700003>.
- [97] Bergmann N, Gyntelberg F, Faber J. The appraisal of chronic stress and the development of the metabolic syndrome: A systematic review of prospective cohort studies. *Endocr Connect.* 2014;**3**:R55–R80. DOI:10.1530/EC-14-0031.
- [98] Siegrist J. Effort—reward imbalance and health in a globalized economy. *Scand J Work Environ Health.* 2008;**suppl 6**:163–168.
- [99] Joksimovic L, Starke D, van der Knesebeck O, Siegrist J. Perceived work stress, overcommitment, and self-reported musculoskeletal pain: A cross-sectional investigation. *Int J Behav Med.* 2002;**9**:122–138.

- [100] Godin I, Kittel F. Differential economic stability and psychosocial stress at work: Associations with psychosomatic complaints and absenteeism. *Soc Sci Med.* 2004;**58**:1543–1553.
- [101] De Jonge J, Bosma H, Peter R, Siegrist J. Job strain, effort-reward imbalance and employee well-being: A large-scale cross-sectional study. *Soc Sci Med.* 2000;**50**:1317–1327.
- [102] Garbarino S, Magnavita N. Work stress and metabolic syndrome in police officers. A prospective study. *PLoS ONE.* 2015;**10**:e0144318. DOI:10.1371/journal.pone.0144318.
- [103] Janczura M, Bochenek G, Nowobilski R, Dropinski J, Kotula-Horowitz K, Laskowicz B, et al. The relationship of metabolic syndrome with stress, coronary heart disease and pulmonary function—An occupational cohort-based study. *PLoS ONE.* 2015;**10**(8):1–20. <http://doi.org/10.1371/journal.pone.0133750>.
- [104] Branth S, Ronquist G, Stridsberg M, Hambræus L, Kindgren E, Olsson R, et al. Development of abdominal fat and incipient metabolic syndrome in young healthy men exposed to long term stress. *Nutr Metab Cardiovasc Dis.* 2007;**17**:427–435.
- [105] Chandola T, Brunner E, Marmot M. Chronic stress at work and the metabolic syndrome: Prospective study. *BMJ (Clinical Research Ed.).* 2006;**332**:521–525.
- [106] Price RH, Choi JN, Vinokur AD. Links in the chain of adversity following job loss: How financial strain and loss of personal control lead to depression, impaired functioning, and poor health. *J Occup Health Psychol.* 2002;**7**:302–312. <http://doi.org/10.1037/1076-8998.7.4.302>.
- [107] Virtanen P, Vahtera J, Kivimäki M, Pentti J, Ferrie J. Employment security and health. *J Epidemiol Community.* 2002;**56**:569–574.
- [108] Dragano N, Verde PE, Siegrist, J. Organisational downsizing and work stress: Testing synergistic health effects in employed men and women. *J Epidemiol Community Health.* 2005;**59**:694–699. <http://doi.org/10.1136/jech.2005.035089>.
- [109] Pyykkonen AJ, Tuomi T, Eriksson JG, Groop L, Isomaa B. Stressful life events and the metabolic syndrome the prevalence, prediction and prevention of diabetes (PPP)-botnia study. *Diabetes Care.* 2010;**33**:378–384.
- [110] Faragher EB, Cass M, Cooper CL. The relationship between job satisfaction and health: A meta-analysis. *Occup Environ Med.* 2005;**62**:105–112.
- [111] Nichole Kelly. Associations between binge eating and executive functioning among young women. [thesis] Richmond VA, US: Virginia Commonwealth University. 2012.
- [112] Dube L, LeBel JL, Lu J. Affect asymmetry and comfort food consumption. *Physiol Behav.* 2005;**86**:559–567. <http://doi.org/10.1016/j.physbeh.2005.08.023>.
- [113] Drescher LS, Hasselbach J. Food Choices under Stress: Considering internet usage and social support. In: AAEA/EAAE/CAES joint symposium social networks, social media and the economics of food Montréal, Québec. Canada, 29–30 May, 2014.

- [114] Ree M, Riediger N, Moghadasian MH. Factors affecting food selection in Canadian population. *Eur J Clin Nutr*. 2008;**62**:1255–1262. <http://doi.org/10.1038/sj.ejcn.1602863>.
- [115] Lampuré A, Schlich P, Deglaire A, Castetbon K, Péneau S, Hercberg S, et al. Sociodemographic, psychological, and lifestyle characteristics are associated with a liking for salty and sweet tastes in French adults. *J Nutr*. 2015;**145**:587–594.
- [116] Camilleri GM, Mejean C, Kesse-Guyot E, Andreeva VA, Bellisle F, Hercberg S, et al. The associations between emotional eating and consumption of energy-dense snack foods are modified by sex and depressive symptomatology. *J Nutr*. 2014;**144**:1264–1273.
- [117] Tan CC, Chow CM. Stress and emotional eating: The mediating role of eating dysregulation. *Pers Individ Differ*. 2014;**66**:1–4. <http://doi.org/10.1016/j.paid.2014.02.033>.
- [118] Oliver G, Wardle J, Gibson EL. Stress and food choice: A laboratory study. *Psychosom Med* 2000;**62**:853–865.
- [119] Tan CC, Ruhl H, Chow CM, Ellis, L. Retrospective reports of parental feeding practices and emotional eating in adulthood: The role of food preoccupation. *Appetite*. 2016;**105**:410–415. DOI:10.1016/j.appet.2016.06.009.
- [120] Eisenstein SA, Bischoff AN, Gredysa DM, Antenor-Dorsey JAV, Koller JM, Al-Lozi A, et al. Emotional eating phenotype is associated with central dopamine d2 receptor binding independent of body mass index. *Sci Rep*. 2015;**5**:11283.
- [121] Park S, Daily JW, Zhang X, Jin HS, Lee HJ, Lee YH. Interactions with the MC4R rs17782313 variant, mental stress and energy intake and the risk of obesity in genome epidemiology study. *Nutr Metab (Lond)*. 2016;**13**:38. DOI:10.1186/s12986-016-0096-8.
- [122] González-Villalva A, Colín-Barenque L, Bizarro-Neves P, Rojas-Lemus M, Rodríguez-Lara V, García-Pelaez I, et al. Pollution by metals: Is there a relationship in glyce-mic control? *Environ Toxicol Pharmacol*. 2016;**46**:337–343. <http://doi.org/10.1016/j.etap.2016.06.023>.
- [123] Cappelletti R, Ceppi M, Claudatus J, Gennaro V. (2016). Health status of male steel workers at an electric arc furnace (EAF) in Trentino, Italy. *J Occup Med Toxicol (Lond)*. 2016;**11**:7. <http://doi.org/10.1186/s12995-016-0095-8>.
- [124] Bahadar H, Mostafalou S, Abdollahi M. Growing burden of diabetes in Pakistan and the possible role of arsenic and pesticides. *J Diabetes Metab Dis*. 2014;**13**:117. DOI: 10.1186/s40200-014-0117-y.
- [125] Bräuner EV, Nordsborg RB, Andersen ZJ, Tjønneland A, Loft S, Raaschou-Nielsen O. Long-term exposure to low-level arsenic in drinking water and diabetes incidence: A prospective study of the diet, cancer and health cohort. *Environ Health Perspect*. 2014;**122**:1059–1065. DOI:10.1289/ehp.1408198.
- [126] Mendez MA, González-Horta C, Sánchez-Ramírez B, Ballinas-Casarrubias L, Hernández Cerón R, Viniegra Morales D, et al. Chronic exposure to arsenic and markers of cardiometabolic risk: A cross-sectional study in Chihuahua, Mexico. *Environ Health Perspect*. 2016;**124**:104–111.

- [127] Moon KA, Guallar E, Umans JG, Devereux RB, Best LG, Francesconi KA, et al. Association between exposure to low to moderate arsenic levels and incident cardiovascular disease. A prospective cohort study. *Ann Intern Med.* 2013;**159**:649–659.
- [128] Jia G, Aroor AR, Martinez-Lemus LA, Sowers JR. Mitochondrial functional impairment in response to environmental toxins in the cardiorenal metabolic syndrome. *Arch Toxicol.* 2015;**89**:147–153.
- [129] Peng S, Liu L, Zhang X, Heinrich J, Zhang J, Schramm KW. et al. A nested case-control study indicating heavy metal residues in meconium associate with maternal gestational diabetes mellitus risk. *Enviro Health.* 2015;**14**:19. DOI: 10.1186/s12940-015-0004-0.
- [130] Feseke SK, Ayotte P, Bouchard M, Levallois P. (Arsenic exposure and type 2 diabetes : Results from the 2007–2009 Canadian health measures survey. *Health Promot Chronic Dis Prev Can.* 2015;**35**:63–72.
- [131] Rahman M, Tondel M, Chowdhury I, Axelson O. Relations between exposure to arsenic, skin lesions, and glucosuria. *Occup Environ Med.* 1999;**56**:277–281.
- [132] Chihuahua ACS, Mendez MA, González-Horta C, Sánchez-Ramírez B, Ballinas-Casarrubias L, Cerón RH, et al. Chronic exposure to arsenic and markers of cardiometabolic risk. *Environ Health Perspect.* 2016;**104**:104–111.
- [133] Peng Q, Harlow SD, Park SK. Urinary arsenic and insulin resistance in US adolescents. *Int J Hyg Environ Health.* 2015;**218**:407–413. doi:10.1016/j.ijheh.2015.03.006.
- [134] Martin E, Gonzalez-Horta C, Rager J, Bailey KA, Sanchez-Ramirez B, Ballinas-Casarrubias L, et al. Metabolomic characteristics of arsenic-associated diabetes in a prospective cohort in Chihuahua, Mexico. *Toxicol Sci.* 2015;**144**:338–346. DOI: 10.1093/toxsci/kfu318.
- [135] Kuo C-C, Howard BV, Umans JG, Gribble MO, Best LG, Francesconi K, et al. Arsenic exposure, arsenic metabolism, and incident diabetes in the strong heart study. *Diabetes Care.* 2015;**38**:620–627.
- [136] Balakrishnan P, Vaidya D, Franceschini N, Voruganti VS, Gribble MO, Haack K, et al. Association of cardiometabolic genes with arsenic metabolism biomarkers in American Indian communities: The strong heart family study (SHFS). *Environ Health Perspect.* 2016 Jun 28. [Epub ahead of print]
- [137] Wang ZX, Jiang CS, Liu L, Wang XH, Jin HJ, Wu Q et al. The role of Akt on arsenic trioxide suppression of 3T3-L1 preadipocyte differentiation. *Cell Res.* 2005;**15**:379–386.
- [138] Divya SP, Pratheeshkumar P, Son YO, Roy RV, Hitron JA, Kim D, et al. Adipocytes and myotubes via oxidative stress-regulated mitochondrial sirt3-FOXO3a signaling pathway. *Toxicol Sci.* 2015;**146**:290–300. DOI:10.1093/toxsci/kfv089.
- [139] Flora SJ. Arsenic-induced oxidative stress and its reversibility. *Free Radic Biol Med.* 2011;**51**:257–281.

- [140] Liu S, Guo X, Wu B, Yu H, Zhang X, Li M. Arsenic induces diabetic effects through beta-cell dysfunction and increased gluconeogenesis in mice. *Sci Rep.* 2014;**4**:6894.
- [141] Simeonova PP, Hulderman T, Harki D, Luster MI. Arsenic exposure accelerates atherogenesis in apolipoprotein E^{-/-} mice. *Environ Health Perspect.* 2003;**111**:1744–1748. DOI: 10.1289/ehp.6332.
- [142] Lemaire M, Negro Silva LF, Lemarié CA, Bolt AM, Flores MOLina M, Krohn RM, et al. Arsenic exposure increases monocyte adhesion to the vascular endothelium, a pro-atherogenic mechanism. *PLoS ONE.* 2015;**10**:e0136592. DOI:10.1371/journal.pone.0136592.
- [143] Kim KN, Park SJ, Choi B, Joo NS. Blood mercury and insulin resistance in nondiabetic Koreans (KNHANES 2008–2010). *Yonsei Med J.* 2015;**56**:944–950. DOI:10.3349/ymj.2015.56.4.944.
- [144] He K, Xun P, Liu K, Morris S, Reis J, Guallar E. Mercury exposure in young adulthood and incidence of diabetes later in life. *Diabetes Care.* Jun 2013;**36**:1584–1589. DOI: 10.2337/dc12-184.
- [145] Moon SS. Association of lead, mercury and cadmium with diabetes in the Korean population: The Korea National Health and Nutrition Examination Survey (KNHANES) 2009–2010. *Diabet Med.* 2013;**30**:e143–e148.
- [146] Kobal AB, Prezelj M, Horvat M, Krsnik M, Gibicar D, Osredkar J. Glutathione level after long-term occupational elemental mercury exposure. *Environ Res.* 2008;**107**:115–123.
- [147] Al-azzawie HF, Umran A, Hyader NH. Oxidative stress, antioxidant status and DNA damage in a mercury exposure workers. *Br J Pharmacol Toxicol.* 2013;**4**:80–88.
- [148] Zeneli L, Sekovanić A, Ajvazi M, Kurti L, Daci N. Alterations in antioxidant defense system of workers chronically exposed to arsenic, cadmium and mercury from coal flying ash. *Environ Geochem Health.* 2016;**38**:65–72. DOI:10.1007/s10653-015-9683-2.
- [149] Tinkov AA, Ajsuvakova OP, Skalnaya MG, Popova EV, Sinitskii AI, Nemereshina ON, et al. Mercury and metabolic syndrome: A review of experimental and clinical observations. *BioMetals.* 2015;**28**:231–254. DOI: 10.1007/s10534-015-9823-2.
- [150] Ya WC, Chun FH, Keh ST, Rong SY, Cheng CY, Ching YY, et al. The role of phosphoinositide 3-kinase/Akt signaling in low-dose mercury-induced mouse pancreatic -cell dysfunction in vitro and in vivo. *Diabetes.* 2006;**55**:1614–1624. DOI: 10.2337/db06-0029.
- [151] Barnes DM, Kircher EA. Effects of mercuric chloride on glucose transport in 3T3-L1 adipocytes. *Toxicol In Vitro.* 2005;**19**:207–214.
- [152] Milnerowicz H, Sciskalska M, Dul M. Pro-inflammatory effects of metals in persons and animals exposed to tobacco smoke. *J Trace Elem Med Biol.* 2015;**29**:1–10. DOI: 10.1016/j.jtemb.2014.04.008.

- [153] Tellez-Plaza M, Navas-Acien A, Caldwell KL, Menke A, Muntner P, Guallar E. Reduction in cadmium exposure in the United States population, 1988–2008: The contribution of declining smoking rates. *Environ Health Perspect.* 2012;**120**:204–209. DOI:10.1289/ehp.1104020.
- [154] Will JC, Galuska DA, Ford ES, Mokdad A, Calle EE. Cigarette smoking and diabetes mellitus: Evidence of a positive association from a large prospective cohort study. *Int J Epidemiol.* 2001;**30**:540–546.
- [155] Sargeant LA, Khaw KT, Bingham S, Day NE, Luben RN, Oakes S, et al. Cigarette smoking and glycaemia: The EPIC-Norfolk study. *European prospective investigation into cancer. Int J Epidemiol.* 2001;**30**:547–554.
- [156] Adhami N, Starck SR, Flores C, Martins Green M. A health threat to bystanders living in the homes of smokers: How smoke toxins deposited on surfaces can cause insulin resistance. *PLoS ONE.* 2016;**11**:e0149510. DOI:10.1371/journal.pone.0149510.
- [157] Afridi HI, Talpur FN, Kazi TG, Brabazon D. Effect of trace and toxic elements of different brands of cigarettes on the essential elemental status of Irish referent and diabetic mellitus consumers. *Biol Trace Elem Res.* 2015;**167**:209. DOI:10.1007/s12011-015-0308-5).
- [158] Yang A, Cheng N, Pu H, Liu S, Dai M, Zheng T, Bai Y. Occupational metal exposures, smoking and risk of diabetes and prediabetes. *Occup Med (Lond).* July 14, 2016., pii. kqw078 [Epub ahead of print].
- [159] Bizoń A, Antonowicz-Juchniewicz J, Milnerowicz M, Śliwińska-Mossoń M, Milnerowicz H. The effect of occupational exposure on pro/antioxidant balance in the blood of non-smoking and smoking smelters with diabetes. *Environ Toxicol Pharmacol.* 2016;**44**:99–106.
- [160] Turyk M, Anderson H, Knobeloch L, Imm P, Persky V. Organochlorine exposure and incidence of diabetes in a cohort of great lakes sport fish consumers. *Environ Health Perspect.* 2009;**117**:1076–1082.
- [161] Pesatori AC, Zocchetti C, Guercilena S, Consonni D, Turrini D, Bertazzi PA. Dioxin exposure and non-malignant health effects: A mortality study. *Occup Environ Med.* 1998;**55**:126–131.
- [162] Vena J, Boffetta P, Becher H, Benn T, Bueno de Mesquita HB, Coggon D, et al. Exposure to dioxin and nonneoplastic mortality in the expanded IARC international cohort study of phenoxy herbicide and chlorophenol production workers and sprayers. *Environ Health Perspect.* 1998;**106**:645–653.
- [163] Shapiro GD, Dodds L, Arbuckle TE, Ashley-Martin J, Fraser W, Fisher M, et al. Exposure to phthalates, bisphenol A and metals in pregnancy and the association with impaired glucose tolerance and gestational diabetes mellitus: The MIREC study. *Environ Int.* 2015;**83**:63–71. DOI: 10.1016/j.envint.2015.05.016.

- [164] García de Aguinaga ML, Posada de la Paz M, Estirado de Cabo E, del Mar Plaza Cano M, Alvarez CS, Saro BB, et al. High prevalence of cardiovascular risk in patients with toxic oil syndrome: A comparative study using the general Spanish population. *Eur J Intern Med.* 2008;**19**:32–39.
- [165] Chang JW, Chen HL, Su HJ, Lee CC. Abdominal obesity and insulin resistance in people exposed to moderate-to-high levels of dioxin. *PLoS ONE.* 2016;**11**:1–14.
- [166] Imbeault P, Findlay CS, Robidoux MA, Haman F, Blais JM, Tremblay A, et al. Dysregulation of cytokine response in Canadian first nations communities: Is there an association with persistent organic pollutant levels? *PLoS ONE.* 2012;**7**:1–8. <http://doi.org/10.1371/journal.pone.0039931>.
- [167] Fujiyoshi PT, Michalek JE, Matsumura F. Molecular epidemiologic evidence for diabetogenic effects of dioxin exposure in U.S. air force veterans of the Vietnam war. *Environ Health Perspect.* 2006;**114**:1677–1683. <http://doi.org/10.1289/ehp.9262>.
- [168] Chang J, Chen H, Su H, Liao P, Guo H, Lee C. Dioxin exposure and insulin resistance in taiwanese living near a highly contaminated area. *Epidemiol.* 2010;**21**:56–61.
- [169] Qiu T, Chen M, Sun X, Cao J, Feng C, Li D, et al. Perfluorooctane sulfonate-induced insulin resistance is mediated by protein kinase B pathway. *Biochem Biophys Res Commun.* 2016;**477**:781–785.
- [170] Seaton A, MacNee W, Donaldson K, Godden D. Particulate air-pollution and acute health-effects. *Lancet.* 1995;**345**:176–178.
- [171] Eze IC, Hemkens LG, Bucher HC, Hoffmann B, Schindler C, Kunzli N, et al. Association between ambient air pollution and diabetes mellitus in Europe and North America: Systematic review and meta-analysis. *Environ Health Perspect.* 2015;**123**:381–389. DOI:10.1289/ehp.1307823.
- [172] Chuang K, Yan Y, Cheng T. Effect of air pollution on blood pressure, blood lipids, and blood sugar: A population-based approach. *J Occup Environ Med.* 2010;**52**:258–262.
- [173] Hansen AB, Ravnskjaer L, Loft S, Andersen KK, Brauner EV, Baastrup R, et al. Long-term exposure to fine particulate matter and incidence of diabetes in the Danish nurse cohort. *Environ Int.* 2016;**91**:243–50.
- [174] Meo SA, Suraya F. Effect of environmental air pollution on cardiovascular diseases. *Eur Rev Med Pharmacol Sci.* 2015;**19**:4890–4897.
- [175] Zheng Z, Xu X, Zhang X, Wang A, Zhang C, Huttermann M, et al. Exposure to ambient particulate matter induces a NASH-like phenotype and impairs hepatic glucose metabolism in an animal model. *J Hepatol.* 2013;**58**:148–154.
- [176] Dennekamp M, Mehenni O, Cherrie JW, Seaton A. Exposure to ultrafine particles and PM_{2.5} in different micro-environments. *Ann Occup Hyg.* 2002;**46**:412–414.

- [177] European Environmental Agency. Emissions of primary PM_{2.5} and PM₁₀ particulate matter, 2014. <http://www.eea.europa.eu/data-and-maps/indicators/emissions-of-primary-particles-and-5/assessment-3>. [Accessed 2016:08:15]
- [178] Shi L, Zanobetti A, Kloog I, Coull BA, Koutrakis P, Melly SJ, et al. Low-concentration PM_{2.5} and mortality: Estimating acute and chronic effects in a population-based study. *Environ Health Perspect*. 2016;**124**:46–52. <http://dx.doi.org/10.1289/ehp.1409111>.
- [179] Pope CA, Turner MC, Burnett RT, Jerrett M, Gapstur SM, Diver WR, et al. Relationships between fine particulate air pollution, cardiometabolic disorders, and cardiovascular mortality. *Circ Res*. 2015;**116**:108–115. <http://doi.org/10.1161/CIRCRESAHA.116.305060>.
- [180] Choi YH, Kim JH, Hong YC. Sex-dependent and body weight-dependent associations between environmental PAHs exposure and insulin resistance: Korean urban elderly panel. *J Epidemiol Commun Health*. 2015;**69**:625–631. DOI:10.1136/jech-2014-204801.
- [181] Kong S, Ding X, Bai Z, et al. A seasonal study of polycyclic aromatic hydrocarbons in PM_{2.5} and PM_{2.5-10} in five typical cities of Liaoning Province, China. *J Hazard Mater*. 2010;**183**:70–80.
- [182] Saldarriaga-Noreña H, López-Márquez R, Murillo-Tovar M, Hernández-Mena L, Ospina-Noreña E, Sánchez-Salinas E, et al. Analysis of PAHs associated with particulate matter PM_{2.5} in two places at the city of Cuernavaca, Morelos, México. *Atmosphere*. 2015;**6**:1259–1270.
- [183] Wei Y, Zhang J, Li Z, Gow A, Chung KF, Hu M, et al. Chronic exposure to air pollution particles increases the risk of obesity and metabolic syndrome: Findings from a natural experiment in Beijing. *FASEB J*. 2016;**30**:2115–2122.
- [184] Khatri M, Bello D, Pal AK, Cohen JM, Woskie S, Gassert T, et al. Evaluation of cytotoxic, genotoxic and inflammatory responses of nanoparticles from photocopiers in three human cell lines. *Part Fibre Toxicol*. 2013;**10**:1–22. <http://doi.org/10.1186/1743-8977-10-42>.
- [185] Brocato J, Hernandez M, Laulicht F, Sun H, Shamy M, Mansour A, et al. In vivo exposures to particulate matter collected from Saudi Arabia or nickel chloride display similar dysregulation of metabolic syndrome genes. *J Toxicol Environ Health Part A*. 2015;**78**:1421–1436. DOI:10.1080/15287394.2015.1095689.
- [186] Qi L, Cornelis MC, Kraft P, Jensen M, van Dam RM, Sun Q, et al. Genetic variants in ABO blood group region, plasma soluble E-selectin levels and risk of type 2 diabetes. *Hum Mol Gen*. 2010;**19**:1856–1862.
- [187] Suadicani P, Hein HO, Gyntelberg F. Airborne occupational exposure, ABO phenotype and risk of ischaemic heart disease in the copenhagen male study. *J Cardiovasc Risk*. 2002;**9**:191–198.
- [188] Wu O, Bayoumi N, Vickers MA, Clark P. ABO(H) blood groups and vascular disease: A systematic review and meta-analysis. *J Thromb Haemos*. 2008;**6**:62–69.

- [189] Kristen E, Cosselman E, Krishnan R. Blood pressure response to controlled diesel exhaust exposure in human subjects. *Hypertension*. 2012;**59**:943–948.
- [190] Devlin RB, Smith CB, Schmitt MT, Rappold AG, Hinderliter A, Graff D, et al. Controlled exposure of humans with metabolic syndrome to concentrated ultrafine ambient particulate matter causes cardiovascular effects. *Toxicol Sci*. 2014;**140**:61–72.
- [191] Baja ES, Schwartz JD, Wellenius GA, Coull BA, Zanobetti A, Vokonas PS, et al. Traffic-related air pollution and QT interval: Modification by diabetes, obesity, and oxidative stress gene polymorphisms in the normative aging study. *Environ Health Perspect*. 2010;**118**:840–846.

INTECH

INTECH